



GCxGC-TOFMS Articles & Reference Materials

- Almstetter, M. F., Oefner, P. J., and Dettmer, K. (2012). Metabolic fingerprinting using comprehensive Two-Dimensional gas chromatography - Time-of-Flight mass spectrometry functional genomics. volume 815 of *Methods in Molecular Biology*, chapter 29, pages 399–411. Springer New York, New York, NY.  
<http://www.springerlink.com/content/jq18821718481614/#section=996908&page=1>
- Antunes, M. M., Lima, S., Fernandes, A., Candeias, J., Pillinger, M., Rocha, S. M., Ribeiro, M. F., and Valente, A. A. (2012). Catalytic dehydration of d-xylose to 2-furfuraldehyde in the presence of Zr-(W,Al) mixed oxides. tracing by-products using two-dimensional gas chromatography-time-of-flight mass spectrometry. *Catalysis Today*.  
<http://www.sciencedirect.com/science/article/pii/S0920586112002325>
- Ari, F., Celikler, S., Oran, S., Balikli, N., Ozturk, S., Ozel, M. Z., Ozyurt, D., and Ulukaya, E. (2012). Genotoxic, cytotoxic, and apoptotic effects of hypogymnia physodes (L.) nyl. on breast cancer cells. *Environ. Toxicol.*, page n/a.  
<http://onlinelibrary.wiley.com/doi/10.1002/tox.21809/abstract;jsessionid=9C4E036764179312E6420903CF18ABAE.d02t03?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Aura, A.-M., Mattila, I., Hyötyläinen, T., Gopalacharyulu, P., Cheynier, V., Souquet, J.-M., Bes, M., Bourvellec, C., Guyot, S., and Orei, M. (2012). Characterization of microbial metabolism of syrah grape products in an in vitro colon model using targeted and non-targeted analytical approaches. *European Journal of Nutrition*, pages 1–14.  
<http://www.springerlink.com/content/104ugw6663651041/>
- Ávila, B. M. F., Vaz, B. G., Pereira, R., Gomes, A. O., Pereira, R. C. L., Corilo, Y. E., Simas, R. C., Nascimento, H. D., Eberlin, M. N., and Azevedo, D. A. (2012). Comprehensive chemical composition of gas oil cuts using Two-Dimensional gas chromatography with Time-of-Flight mass spectrometry and electrospray ionization coupled to fourier transform ion cyclotron resonance mass spectrometry. *Energy Fuels*, 26(8):5069–5079. <http://pubs.acs.org/doi/abs/10.1021/ef300631e>
- Baidoo, E. E. K., Benke, P. I., and Keasling, J. D. (2012). Mass Spectrometry-Based microbial metabolomics microbial systems biology. volume 881 of *Methods in Molecular Biology*, chapter 9, pages 215–278. Humana Press, Totowa, NJ.  
<http://www.springerlink.com/content/k6k2j5368r646h33/#section=1075588&page=1>
- Bastos, P. M. and Haglund, P. (2012). The use of comprehensive two-dimensional gas chromatography and structure-activity modeling for screening and preliminary risk assessment of organic contaminants in soil, sediment, and surface water. *Journal of Soils and Sediments*, 12(7):1079–1088. <http://www.springerlink.com/content/q7x52qk712161146/>
- Bean, H. D., Dimandja, J.-M. D., and Hill, J. E. (2012). Bacterial volatile discovery using solid phase microextraction and comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry. *Journal of Chromatography B*, 901:41–46.  
<http://www.sciencedirect.com/science/article/pii/S1570023212003194>
- Beckstrom, A. C., Tanya, P., Humston, E. M., Snyder, L. R., Synovec, R. E., and Juul, S. E. (2012). The perinatal transition of the circulating metabolome in a nonhuman primate. *Pediatric Research*, 71(4-1):338–344. <http://www.nature.com/pr/journal/v71/n4-1/full/pr201174a.html>
- Boogaard, P. J., Goyak, K. O., Biles, R. W., van Stee, L. L. P., Miller, M. S., and Miller, M. J. (2012). Comparative toxicokinetics of low-viscosity mineral oil in fischer 344 rats, Sprague-Dawley rats, and humans - implications for an acceptable daily intake (ADI). *Regulatory Toxicology and Pharmacology*, 63(1):69–77. <http://www.sciencedirect.com/science/article/pii/S0273230012000402>
- Brasseur, C., Dekeirsschietier, J., Schotsmans, E. M. J., de Koning, S., Wilson, A. S., Haubruge, E., and Focant, J.-F. (2012). Comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry for the forensic study of cadaveric volatile organic compounds released in soil by buried decaying pig carcasses. *Journal of Chromatography A*, 1255:163–170.  
<http://www.sciencedirect.com/science/article/pii/S002196731200461X>
- Caldeira, M., Perestrelo, R., Barros, A. S., Bilelo, M. J., Morête, A., Cãmara, J. S., and Rocha, S. M. (2012). Allergic asthma exhaled breath metabolome: A challenge for comprehensive two-dimensional gas chromatography. *Journal of Chromatography A*, 1254:87–97.  
<http://www.sciencedirect.com/science/article/pii/S0021967312010576>
- Cao, G., Cai, H., Cong, X., Liu, X., Ma, X., Lou, Y., Qin, K., and Cai, B. (2012). Global detection and analysis of volatile components from sun-dried and sulfur-fumigated herbal medicine by comprehensive two-dimensional gas chromatography/time-of-flight mass spectrometry. *Analyst*, 137(16):3828–3835. <http://pubs.rsc.org/en/Content/ArticleLanding/2012/AN/c2an35543d>
- Carmichael, C. A., Arey, J. S., Graham, W. M., Linn, L. J., Lemkau, K. L., Nelson, R. K., and Reddy, C. M. (2012). Floating oil-covered debris from deepwater horizon: identification and application. *Environmental Research Letters*, 7(1):015301+.  
<http://iopscience.iop.org/1748-9326/7/1/015301/>
- de la Mata, A. P. and Harynuk, J. J. (2012). Limits of detection and quantification in comprehensive multidimensional separations. 1. a theoretical look. *Anal. Chem.*, 84(15):6646–6653. <http://pubs.acs.org/doi/abs/10.1021/ac3010204>
- Dekeirsschietier, J., Stefanuto, P.-H., Brasseur, C., Haubruge, E., and Focant, J.-F. (2012). Enhanced characterization of the smell of death by comprehensive Two-Dimensional gas Chromatography-Time-of-Flight mass spectrometry (GCxGC-TOFMS). *PLoS ONE*, 7(6):e39005+.  
<http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0039005>
- Drabova, L., Pulkrabova, J., Kalachova, K., Tomaniova, M., Kocourek, V., and Hajslova, J. (2012). Rapid determination of polycyclic aromatic hydrocarbons (PAHs) in tea using two-dimensional gas chromatography coupled with time of flight mass spectrometry. *Talanta*, 100:207–216. <http://www.sciencedirect.com/science/article/pii/S0039914012006704>

- Dutriez, T., Courtiade, M., Ponthus, J., Thiébaud, D., Dulot, H., and Hennion, M.-C. (2012). Complementarity of fourier transform ion cyclotron resonance mass spectrometry and high temperature comprehensive two-dimensional gas chromatography for the characterization of resin fractions from vacuum gas oils. *Fuel*, 96:108–119. <http://www.sciencedirect.com/science/article/pii/S0016236111007848>
- Eiserbeck, C., Nelson, R. K., Grice, K., Curiale, J., and Reddy, C. M. (2012). Comparison of GC-MS, GC-MRM-MS, and GCxGC to characterise higher plant biomarkers in tertiary oils and rock extracts. *Geochimica et Cosmochimica Acta*, 87:299–322. <http://www.sciencedirect.com/science/article/pii/S0016703712001858>
- Gauchotte-Lindsay, C., Richards, P., McGregor, L. A., Thomas, R., and Kalin, R. M. (2012). A one-step method for priority compounds of concern in tar from former industrial sites: Trimethylsilyl derivatisation with comprehensive two-dimensional gas chromatography. *Journal of Chromatography A*, 1253:154–163. <http://www.sciencedirect.com/science/article/pii/S0021967312010308>
- Hanus, R., obotník, J., Krasulová, J., Jiro, P., áek, P., Kalinová, B., Dolejová, K., Cvaka, J., Bourguignon, T., Roisin, Y., Lacey, M. J., and Sillam-Dussès, D. (2012). Nonadecadienone, a new termite Trail-Following pheromone identified in *glossotermes oculatus* (serritermitidae). *Chemical Senses*, 37(1):55–63. <http://chemse.oxfordjournals.org/content/37/1/55>
- Harvey, S. D. and Wahl, J. H. (2012). On-matrix derivatization for dynamic headspace sampling of nonvolatile surface residues. *Journal of Chromatography A*, 1256:58–66. <http://www.sciencedirect.com/science/article/pii/S0021967312011879>
- Hoh, E., Dodder, N. G., Lehotay, S. J., Pangallo, K. C., Reddy, C. M., and Maruya, K. A. (2012). Nontargeted comprehensive Two-Dimensional gas Chromatography/Time-of-Flight mass spectrometry method and software for inventorying persistent and bioaccumulative contaminants in marine environments. *Environ. Sci. Technol.*, 46(15):8001–8008. <http://pubs.acs.org/doi/abs/10.1021/es301139q>
- Hyotylainen, T., Mattila, I., Wiedmer, S. K., Koivuniemi, A., Taskinen, M.-R., Yki-Jarvinen, H., and Oresic, M. (2012). Metabolomic analysis of polar metabolites in lipoprotein fractions identifies lipoprotein-specific metabolic profiles and their association with insulin resistance. *Mol. BioSyst.*, 8(10):2559–2565. <http://pubs.rsc.org/en/Content/ArticleLanding/2012/MB/c2mb25115a>
- Jalali, H. T., Petronilho, S., Villaverde, J. J., Coimbra, M. A., Domingues, Ebrahimian, Z. J., Silvestre, A. J. D., and Rocha, S. M. (2012). Deeper insight into the monoterpene composition of *ferula gummosa* oleo-gum-resin from iran. *Industrial Crops and Products*, 36(1):500–507. <http://www.sciencedirect.com/science/article/pii/S0926669011004262>
- Kalachova, K., Pulkrabova, J., Cajka, T., Drabova, L., and Hajslova, J. (2012). Implementation of comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry for the simultaneous determination of halogenated contaminants and polycyclic aromatic hydrocarbons in fish. *Analytical and Bioanalytical Chemistry*, 403(10):2813–2824. <http://www.springerlink.com/content/v563k72661631ul/>
- Kiefl, J., Cordero, C., Nicolotti, L., Schieberle, P., Reichenbach, S. E., and Bicchi, C. (2012). Performance evaluation of non-targeted peak-based cross-sample analysis for comprehensive two-dimensional gas chromatography-mass spectrometry data and application to processed hazelnut profiling. *Journal of Chromatography A*, 1243:81–90. <http://www.sciencedirect.com/science/article/pii/S0021967312006401>
- Koo, I., Zhao, Y., Zhang, J., Kim, S., and Zhang, X. (2012). A method of calculating the second dimension hold-up time for comprehensive two-dimensional gas chromatography. *Journal of Chromatography A*. <http://www.sciencedirect.com/science/article/pii/S0021967312013027>
- Lee, J. Y., Lane, D. A., Heo, J. B., Yi, S.-M., and Kim, Y. P. (2012). Quantification and seasonal pattern of atmospheric reaction products of gas phase PAHs in PM2.5. *Atmospheric Environment*, 55:17–25. <http://www.sciencedirect.com/science/article/pii/S1352231012002397>
- Li, S., Hu, S., Cao, J., Wu, M., and Zhang, D. (2012). Diamondoid characterization in condensate by comprehensive Two-Dimensional gas chromatography with Time-of-Flight mass spectrometry: The junggar basin of northwest china. *International Journal of Molecular Sciences*, 13(9):11399–11410. <http://www.mdpi.com/1422-0067/13/9/11399>
- Livngood, P., Maciejewski, R., Chen, W., and Ebert, D. (2012). OmicsVis: an interactive tool for visually analyzing metabolomics data. *BMC Bioinformatics*, 13(Suppl 8):S6+. <http://www.ncbi.nlm.nih.gov/pubmed/22607515>
- Maity, S., Jannasch, A., Adamec, J., Nalepa, T., Höök, T. O., and Sepúlveda, M. S. (2012). Starvation causes disturbance in amino acid and fatty acid metabolism in *diporeia*. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, 161(4):348–355. <http://www.sciencedirect.com/science/article/pii/S109649591200005X>
- Mal, M., Koh, P. K., Cheah, P. Y., and Chan, E. C. (2012). Metabotyping of human colorectal cancer using two-dimensional gas chromatography mass spectrometry. *Analytical and Bioanalytical Chemistry*, 403(2):483–493. <http://www.springerlink.com/content/k42667t0w5313225/>
- Malleswara Rao, T. V., Dupain, X., and Makkee, M. (2012). Fluid catalytic cracking: Processing opportunities for Fischer-Tropsch waxes and vegetable oils to produce transportation fuels and light olefins. *Microporous and Mesoporous Materials*. <http://www.sciencedirect.com/science/article/pii/S1387181112004362>
- Manzano, C., Hoh, E., and Simonich, S. L. (2012). Improved separation of complex polycyclic aromatic hydrocarbon mixtures using novel column combinations in GC x GC/ToF-MS. *Environ. Sci. Technol.*, 46(14):7677–7684. <http://pubs.acs.org/doi/abs/10.1021/es301790h>
- McGinitie, T. M. and Harynyk, J. J. (2012). Prediction of retention times in comprehensive two-dimensional gas chromatography using thermodynamic models. *Journal of Chromatography A*, 1255:184–189. <http://www.sciencedirect.com/science/article/pii/S002196731200266X>

- McGregor, L. A., Gauchotte-Lindsay, C., Nic Daéid, N., Thomas, R., and Kalin, R. M. (2012). Multivariate statistical methods for the environmental forensic classification of coal tars from former manufactured gas plants. *Environ. Sci. Technol.*, 46(7):3744–3752. <http://pubs.acs.org/doi/abs/10.1021/es203708w>
- Meinert, C., Filippi, J.-J., deã...Marcellus, P., Leã...Sergeantã...d'Hendecourt, L., and Meierhenrich, U. J. (2012). N-(2-Aminoethyl)glycine and amino acids from interstellar ice analogues. *ChemPlusChem*, 77(3):186–191. <http://onlinelibrary.wiley.com/doi/10.1002/cplu.201100048/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Meinert, C. and Meierhenrich, U. J. (2012). A new dimension in separation science: Comprehensive Two-Dimensional gas chromatography. *Angew. Chem. Int. Ed.*, page n/a. <http://onlinelibrary.wiley.com/doi/10.1002/anie.201200842/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Menéndez-Carreño, M., Steenbergen, H., and Janssen, H.-G. (2012). Development and validation of a comprehensive two-dimensional gas chromatography-mass spectrometry method for the analysis of phytosterol oxidation products in human plasma. *Analytical and Bioanalytical Chemistry*, 402(6):2023–2032. <http://www.springerlink.com/content/x03717977g56250m/>
- Moraes, M. S., Georges, F., Almeida, S. R., Damasceno, F. C., Maciel, G. P., Zini, C. A., Jacques, R. A., and Caramão, E. B. (2012a). Analysis of products from pyrolysis of brazilian sugar cane straw. *Fuel Processing Technology*, 101:35–43. <http://www.sciencedirect.com/science/article/pii/S0378382012000975>
- Moraes, M. S., Migliorini, M. V., Damasceno, F. C., Georges, F., Almeida, S., Zini, C. A., Jacques, R. A., and Caramão, E. B. (2012b). Qualitative analysis of bio oils of agricultural residues obtained through pyrolysis using comprehensive two dimensional gas chromatography with time-of-flight mass spectrometric detector. *Journal of Analytical and Applied Pyrolysis*. <http://www.sciencedirect.com/science/article/pii/S016523701200099X>
- Mostafa, A., Edwards, M., and Górecki, T. (2012). Optimization aspects of comprehensive two-dimensional gas chromatography. *Journal of Chromatography A*, 1255:38–55. <http://www.sciencedirect.com/science/article/pii/S0021967312003706>
- Munoz, M., de Pedro, Z. M., Casas, J. A., and Rodriguez, J. J. (2012). Triclosan breakdown by fenton-like oxidation. *Chemical Engineering Journal*, 198-199:275–281. <http://www.sciencedirect.com/science/article/pii/S1385894712006870>
- Muusse, M., Langford, K., Tollefsen, K. E., Cornelissen, G., Haglund, P., Hylland, K., and Thomas, K. V. (2012). Characterization of AhR agonist compounds in roadside snow. *Analytical and Bioanalytical Chemistry*, 403(7):2047–2056. <http://www.springerlink.com/content/v6768t2u37127802/>
- Naudé, Y. and Rohwer, E. R. (2012). Two multidimensional chromatographic methods for enantiomeric analysis of o,p'-DDT and o,p'-DDD in contaminated soil and air in a malaria area of south africa. *Analytica Chimica Acta*, 730:120–126. <http://www.sciencedirect.com/science/article/pii/S000326701200445X>
- Ngo, H. L., Hoh, E., and Foglia, T. A. (2012). Improved synthesis and characterization of saturated branched-chain fatty acid isomers. *Eur. J. Lipid Sci. Technol.*, 114(2):213–221. <http://onlinelibrary.wiley.com/doi/10.1002/ejlt.201000471/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Nizio, K. D. and Harynyuk, J. J. (2012). Analysis of alkyl phosphates in petroleum samples by comprehensive two-dimensional gas chromatography with nitrogen phosphorus detection and post-column deans switching. *Journal of Chromatography A*, 1252:171–176. <http://www.sciencedirect.com/science/article/pii/S0021967312009648>
- Oliveira, C. R., Ferreira, A. A., Oliveira, C. J. F., Azevedo, D. A., Santos Neto, E. V., and Aquino Neto, F. R. (2012a). Biomarkers in crude oil revealed by comprehensive two-dimensional gas chromatography time-of-flight mass spectrometry: Depositional paleoenvironment proxies. *Organic Geochemistry*, 46:154–164. <http://www.sciencedirect.com/science/article/pii/S0146638012000526>
- Oliveira, C. R., Oliveira, C. J. F., Ferreira, A. A., Azevedo, D. A., and Aquino Neto, F. R. (2012b). Characterization of aromatic steroids and hopanoids in marine and lacustrine crude oils using comprehensive two dimensional gas chromatography coupled to time-of-flight mass spectrometry (GCxGC-TOFMS). *Organic Geochemistry*. <http://www.sciencedirect.com/science/article/pii/S0146638012001246>
- Oliveira, V., Santos, A. L., Aguiar, C., Santos, L., Salvador, A. C., Gomes, N. C. M., Silva, H., Rocha, S. M., Almeida, A., and Cunha, A. (2012c). Prokaryotes in salt marsh sediments of ria de aveiro: Effects of halophyte vegetation on abundance and diversity. *Estuarine, Coastal and Shelf Science*, 110:61–68. <http://www.sciencedirect.com/science/article/pii/S0272771412000868>
- Paulina de la Mata, A., Nizio, K. D., and Harynyuk, J. J. (2012). Integration parameters and their effects on quantitative results with two-step peak summation quantitation in comprehensive two-dimensional gas chromatography. *Journal of Chromatography A*, 1255:190–195. <http://www.sciencedirect.com/science/article/pii/S0021967312003196>
- Payeur, A. L., Lorenz, M. A., and Kennedy, R. T. (2012). Analysis of fatty acid composition in insulin secreting cells by comprehensive two-dimensional gas chromatography time-of-flight mass spectrometry. *Journal of Chromatography B*, 893-894:187–192. <http://www.sciencedirect.com/science/article/pii/S1570023212001523>
- Pena-Abaurrea, M., Ye, F., Blasco, J., and Ramos, L. (2012). Evaluation of comprehensive two-dimensional gas chromatography-time-of-flight-mass spectrometry for the analysis of polycyclic aromatic hydrocarbons in sediments. *Journal of Chromatography A*, 1256:222–231. <http://www.sciencedirect.com/science/article/pii/S0021967312011156>



- Risticvic, S., DeEll, J. R., and Pawlisygn, J. (2012). Solid phase microextraction coupled with comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry for high-resolution metabolite profiling in apples: Implementation of structured separations for optimization of sample preparation procedure in complex samples. *Journal of Chromatography A*, 1251:208–218. <http://www.sciencedirect.com/science/article/pii/S0021967312009466>
- Rocha, S. M., Caldeira, M., Carrola, J., Santos, M., Cruz, N., and Duarte, I. F. (2012). Exploring the human urine metabolomic potentialities by comprehensive two-dimensional gas chromatography coupled to time of flight mass spectrometry. *Journal of Chromatography A*, 1252:155–163. <http://www.sciencedirect.com/science/article/pii/S0021967312009612>
- Rosenstiel, T. N., Shortlidge, E. E., Melnychenko, A. N., Pankow, J. F., and Eppley, S. M. (2012). Sex-specific volatile compounds influence microarthropod-mediated fertilization of moss. *Nature*, 489(7416):431–433. <http://www.deepdyve.com/lp/nature/sex-specific-volatile-compounds-influence-microarthropod-mediated-q1rKM0rkCL?key=citeulike>
- Sanda, M., áek, P., Streinz, L., Draínský, M., and Koutek, B. (2012). Profiling and characterization of volatile secretions from the european stink bug *graphosoma lineatum* (heteroptera: Pentatomidae) by two-dimensional gas chromatography/time-of-flight mass spectrometry. *Journal of Chromatography B*, 881–882:69–75. <http://www.sciencedirect.com/science/article/pii/S1570023211007847>
- Santiago-Morales, J., Gómez, M. J., Herrera, S., Fernández-Alba, A. R., García-Calvo, E., and Rosal, R. (2012). Oxidative and photochemical processes for the removal of galaxolide and tonalide from wastewater. *Water Research*, 46(14):4435–4447. <http://www.sciencedirect.com/science/article/pii/S0043135412003892>
- Schäffer, M., Gröger, T., Pütz, M., Dieckmann, S., and Zimmermann, R. (2012). Comparative analysis of the chemical profiles of 3,4-Methylenedioxyamphetamine based on comprehensive Two-Dimensional gas Chromatography-Time-of-Flight mass spectrometry (GCxGC-TOFMS)\*. *Journal of Forensic Sciences*, 57(5):1181–1189. <http://onlinelibrary.wiley.com/doi/10.1111/j.1556-4029.2012.02137.x/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Schoeman, J. C., du Preez, I., and Loots, D. T. (2012). A comparison of four sputum pre-extraction preparation methods for identifying and characterising mycobacterium tuberculosis using GCxGC-TOFMS metabolomics. *Journal of Microbiological Methods*. [ScienceDirect.com - Journal of Microbiological Methods - A comparison of four sputum pre-extraction preparation methods for identifying and characterising Mycobacterium tuberculosis using GCxGC-TOFMS metabolomics](http://www.sciencedirect.com/science/article/pii/S0168165612000389)
- Stadler, S., Stefanuto, P.-H., Byer, J. D., Brokl, M., Forbes, S., and Focant, J.-F. (2012). Analysis of synthetic canine training aids by comprehensive two-dimensional gas chromatography-time of flight mass spectrometry. *Journal of Chromatography A*, 1255:202–206. <http://www.sciencedirect.com/science/article/pii/S0021967312005389>
- Tcaciuc, A. P., Nelson, R. K., Reddy, C. M., and Gschwend, P. M. (2012). Estimating phospholipid Membrane-Water partition coefficients using comprehensive Two-Dimensional gas chromatography. *Environ. Sci. Technol.*, 46(6):3449–3456. <http://pubs.acs.org/doi/abs/10.1021/es203792e>
- Tissot, E., Rochat, S., Debonneville, C., and Chaintreau, A. (2012). Rapid GC-FID quantification technique without authentic samples using predicted response factors. *Flavour Fragr. J.*, 27(4):290–296. <http://onlinelibrary.wiley.com/doi/10.1002/ffj.3098/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Vaníková, L., Svato, A., Kroiss, J., Kaltenpoth, M., Nascimento, R. R., Hoskovec, M., Bizová, R., and Kalinová, B. (2012). Cuticular hydrocarbons of the south american fruit fly *Anastrepha fraterculus*: Variability with sex and age. *Journal of Chemical Ecology*, pages 1–10. <http://www.springerlink.com/content/4172400512504241/>
- Ventura, G. T., Simoneit, B. R. T., Nelson, R. K., and Reddy, C. M. (2012). The composition, origin and fate of complex mixtures in the maltene fractions of hydrothermal petroleum assessed by comprehensive two-dimensional gas chromatography. *Organic Geochemistry*, 45:48–65. <http://www.sciencedirect.com/science/article/pii/S0146638012000034>
- Villière, A., Arvisenet, G., Lethuaut, L., Prost, C., and Sérot, T. (2012). Selection of a representative extraction method for the analysis of odourant volatile composition of french cider by GC-MS-O and GCxGC-TOF-MS. *Food Chemistry*, 131(4):1561–1568. <http://www.sciencedirect.com/science/article/pii/S0308814611014403>
- Vivó-Truyols, G. (2012). Bayesian approach for peak detection in Two-Dimensional chromatography. *Anal. Chem.*, 84(6):2622–2630. <http://pubs.acs.org/doi/abs/10.1021/ac202124t>
- Wang, H., Zhang, S., Weng, N., Wei, X., Zhu, G., Yu, H., Bi, L., and Ma, W. (2012a). Insight of unresolved complex mixtures of saturated hydrocarbons in heavy oil via GCxGC-TOFMS analysis. *SCIENCE CHINA Chemistry*, pages 1–9. <http://www.springerlink.com/content/k411780654m13rk6/>
- Wang, K., Zhu, R.-z., Qu, R.-f., and Li, Z.-y. (2012b). Comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry for the analysis of volatile components in neroli essential oil. *Mendeleev Communications*, 22(1):45–46. <http://www.sciencedirect.com/science/article/pii/S0959943612000193>
- Wang, X., Li, P., Zhang, W., Zhang, Q., Ma, F., Yu, L., and Wang, L. (2012). Screening for pesticide residues in oil seeds using solid-phase dispersion extraction and comprehensive two-dimensional gas chromatography time-of-flight mass spectrometry. *J. Sep. Science*, 35(13):1634–1643. <http://onlinelibrary.wiley.com/doi/10.1002/jssc.201100863/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Wei, X., Shi, X., Kim, S., McClain, C., and Zhang, X. (2012). A novel Two-Stage alignment method for liquid chromatography mass Spectrometry-Based metabolomics intelligent computing technology. volume 7389 of *Lecture Notes in Computer Science*, chapter 20, pages 152–159. Springer Berlin / Heidelberg, Berlin, Heidelberg. <http://www.springerlink.com/content/t080t1v63xk21443/>

- Welke, J. E., Manfroi, V., Zanus, M., Lazarotto, M., and Alcaraz Zini, C. (2012). Characterization of the volatile profile of brazilian merlot wines through comprehensive two dimensional gas chromatography time-of-flight mass spectrometric detection. *Journal of Chromatography A*, 1226:124–139. <http://www.sciencedirect.com/science/article/pii/S0021967312000349>
- Weusten, J. J. A. M., Derks, E. P. P. A., Mommers, J. H. M., and van der Wal, S. (2012). Alignment and clustering strategies for GCxGC-MS features using a cylindrical mapping. *Analytica Chimica Acta*, 726:9–21. <http://www.sciencedirect.com/science/article/pii/S0003267012003741>
- Ye, G., Zhu, B., Yao, Z., Yin, P., Lu, X., Kong, H., Fan, F., Jiao, B., and Xu, G. (2012). Analysis of urinary metabolic signatures of early hepatocellular carcinoma recurrence after surgical removal using gas Chromatography–Mass spectrometry. *J. Proteome Res.*, 11(8):4361–4372. <http://pubs.acs.org/doi/abs/10.1021/pr300502v>
- Zeng, Z.-D., Hugel, H. M., and Marriott, P. J. (2012). Component correlation between related samples by using comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry with chemometric tools. *Journal of Chromatography A*, 1254:98–106. <http://www.sciencedirect.com/science/article/pii/S0021967312010679>
- Zhang, C., Eganhouse, R. P., Pontolillo, J., Cozzarelli, I. M., and Wang, Y. (2012). Determination of nonylphenol isomers in landfill leachate and municipal wastewater using steam distillation extraction coupled with comprehensive two-dimensional gas chromatography/time-of-flight mass spectrometry. *Journal of Chromatography A*, 1230:110–116. <http://www.sciencedirect.com/science/article/pii/S0021967312000714>
- Zhu, S., He, S., Worton, D. R., and Goldstein, A. H. (2012). Predictions of comprehensive two-dimensional gas chromatography separations from isothermal data. *Journal of Chromatography A*, 1233:147–151. <http://www.sciencedirect.com/science/article/pii/S0021967312002919>
- Zubot, W., MacKinnon, M. D., Chelme-Ayala, P., Smith, D. W., and Gamal El-Din, M. (2012). Petroleum coke adsorption as a water management option for oil sands process-affected water. *Science of The Total Environment*, 427-428:364–372. <http://www.sciencedirect.com/science/article/pii/S0048969712005219>

## 2011

---

- Almstetter, M. F., Appel, I. J., Dettmer, K., Gruber, M. A., and Oefner, P. J. (2011). Comparison of two algorithmic data processing strategies for metabolic fingerprinting by comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry. *Journal of Chromatography A*. <http://www.sciencedirect.com/science/article/pii/S002196731101171X>
- Ávila, B. M. F., Pereira, R., Gomes, A. O., and Azevedo, D. A. (2011). Chemical characterization of aromatic compounds in extra heavy gas oil by comprehensive two-dimensional gas chromatography coupled to time-of-flight mass spectrometry. *Journal of Chromatography A*, 1218(21):3208–3216. <http://www.sciencedirect.com/science/article/pii/S0021967310012872>
- Beckstrom, A. C., Humston, E. M., Snyder, L. R., Synovec, R. E., and Juul, S. E. (2011). Application of comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometry method to identify potential biomarkers of perinatal asphyxia in a non-human primate model. *Journal of Chromatography A*, 1218(14):1899–1906. <http://www.sciencedirect.com/science/article/pii/S0021967311001828>
- Bertrand, E., Machado-Maturana, E., Chevarin, C., Portanguen, S., Mercier, F., Tournayre, P., Abouelkaram, S., Guillard, A.-S., Kondjoyan, A., and Berdagué, J.-L. (2011). Heat-induced volatiles and odour-active compounds in a model cheese. *International Dairy Journal*, 21(10):806–814. <http://www.sciencedirect.com/science/article/pii/S0958694611001014>
- Breet, E., Nortjé, Y., and van Greuning, C. (2011). Supercritical carbon dioxide extracted oil from *jatropha curcas*: Directive for the biodiesel industry? *The Journal of Supercritical Fluids*, 60:38–44. <http://www.sciencedirect.com/science/article/pii/S0896844611003081>
- Cheong, K. W., Tan, C. P., Mirhosseini, H., Chin, S. T., Che Man, Y. B., Hamid, N. S., Osman, A., and Basri, M. (2011). Optimization of equilibrium headspace analysis of volatile flavor compounds of malaysian soursop (*annona muricata*): Comprehensive two-dimensional gas chromatography time-of-flight mass spectrometry (GCxGC-TOFMS). *Food Chemistry*, 125(4):1481–1489. <http://www.sciencedirect.com/science/article/pii/S0308814610013245>
- Chin, S.-T., Eyres, G. T., and Marriott, P. J. (2011). Identification of potent odourants in wine and brewed coffee using gas chromatography-olfactometry and comprehensive two-dimensional gas chromatography. *Journal of Chromatography A*, 1218(42):7487–7498. <http://www.sciencedirect.com/science/article/pii/S002196731100851X>
- da Silva, J. M., Zini, C. A., and Caramão, E. B. (2011). Evaluation of comprehensive two-dimensional gas chromatography with micro-electron capture detection for the analysis of seven pesticides in sediment samples. *Journal of Chromatography A*, 1218(21):3166–3172. <http://www.sciencedirect.com/science/article/pii/S0021967310016870>
- de Marcellus, P., Meinert, C., Nuevo, M., Filippi, J.-J., Danger, G., Deboffle, D., Nahon, L., d'Hendecourt, L. L. S., and Meierhenrich, U. J. (2011). Non-racemic amino acid production by ultraviolet irradiation of achiral interstellar ice analogs with circularly polarized light. *The Astrophysical Journal Letters*, 727(2):L27+. <http://iopscience.iop.org/2041-8205/727/2/L27/>
- de Vos, J., Dixon, R., Vermeulen, G., Gorst-Allman, P., Cochran, J., Rohwer, E., and Focant, J.-F. (2011a). Comprehensive two-dimensional gas chromatography time of flight mass spectrometry (GCxGC-TOFMS) for environmental forensic investigations in developing countries. *Chemosphere*, 82(9):1230–1239. <http://www.sciencedirect.com/science/article/pii/S004565351001444X>
- de Vos, J., Gorst-Allman, P., and Rohwer, E. (2011b). Establishing an alternative method for the quantitative analysis of polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans by comprehensive two dimensional gas chromatography-time-of-flight mass spectrometry for developing countries. *Journal of Chromatography A*, 1218(21):3282–3290. <http://www.sciencedirect.com/science/article/pii/S0021967311004134>

- Edwards, M., Mostafa, A., and Górecki, T. (2011). Modulation in comprehensive two-dimensional gas chromatography: 20 years of innovation. *Analytical and Bioanalytical Chemistry*, 401(8):2335–2349. <http://www.springerlink.com/content/b3p0884055h216t2/>
- Eiserbeck, C., Nelson, R. K., Grice, K., Curiale, J., Reddy, C. M., and Raiteri, P. (2011). Separation of 18 $\alpha$ (H)-, 18 $\beta$ (H)-oleanane and lupane by comprehensive two-dimensional gas chromatography. *Journal of Chromatography A*, 1218(32):5549–5553. <http://www.sciencedirect.com/science/article/pii/S0021967311008338>
- Fraga, C. G., Wahl, J. H., and Núñez, S. P. (2011). Profiling of volatile impurities in tetramethylenedisulfotetramine (TETS) for synthetic-route determination. *Forensic Science International*, 210(1-3):164–169. [http://www.fsijournal.org/article/S0379-0738\(11\)00111-3/abstract](http://www.fsijournal.org/article/S0379-0738(11)00111-3/abstract)
- Gardner, J. Y., Brillhart, D. E., Benjamin, M. M., Dixon, L. G., Mitchell, L. M., and Dimandja, J.-M. D. (2011). The use of GCxGC/TOF MS with multivariate analysis for the characterization of foodborne pathogen bacteria profiles. *J. Sep. Science*, 34(2):176–185. <http://onlinelibrary.wiley.com/doi/10.1002/jssc.201000612/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Göü, F., Lewis, A. C., and Özel, M. Z. (2011). Analysis of black mulberry volatiles using GCxGC-TOF/MS. *International Journal of Food Properties*, 14(1):29–36. <http://www.tandfonline.com/doi/abs/10.1080/10942910903125276>
- Gogus, F., Özel, M. Z., Kocak, D., Hamilton, J. F., and Lewis, A. C. (2011). Analysis of roasted and unroasted pistacia terebinthus volatiles using direct thermal desorption-GCxGC-TOF/MS. *Food Chemistry*, 129(3):1258–1264. <http://www.sciencedirect.com/science/article/pii/S0308814611006789>
- Götz, A., Tyynismaa, H., Euro, L., Ellonen, P., Hyötyläinen, T., Ojala, T., Hämäläinen, R. H., Tommiska, J., Raivio, T., Oresic, M., Karikoski, R., Tammela, O., Simola, K. O. J., Paetau, A., Tyni, T., and Suomalainen, A. (2011). Exome sequencing identifies mitochondrial Alanyl-tRNA synthetase mutations in infantile mitochondrial cardiomyopathy. *The American Journal of Human Genetics*, 88(5):635–642. <http://www.cell.com/AJHG/retrieve/pii/S0002929711001479>
- Gröger, T. and Zimmermann, R. (2011). Application of parallel computing to speed up chemometrics for GCxGC-TOFMS based metabolic fingerprinting. *Talanta*, 83(4):1289–1294. <http://www.sciencedirect.com/science/article/pii/S0039914010007101>
- Ha, J., Seo, D., and Shin, D. (2011). Determination of elaidic and vaccenic acids in foods using GCxGC-FID and GCxGC-TOFMS. *Talanta*, 85(1):252–258. <http://www.sciencedirect.com/science/article/pii/S003991401100258X>
- Harynuk, J. J., Rossé, A. D., and McGarvey, G. B. (2011). Study of alkyl phosphates in industrial petroleum mixtures by comprehensive two-dimensional gas chromatography time-of-flight mass spectrometry. *Analytical and Bioanalytical Chemistry*, 401(8):2415–2422. <http://www.springerlink.com/content/a610524028q12h36/>
- Johanningsmeier, S. D. and McFeeters, R. F. (2011). Detection of volatile spoilage metabolites in fermented cucumbers using nontargeted, comprehensive 2-Dimensional gas Chromatography-Time-of-Flight mass spectrometry (GCxGC-TOFMS). *Journal of Food Science*, 76(1):C168–C177. <http://onlinelibrary.wiley.com/doi/10.1111/j.1750-3841.2010.01918.x/abstract?jsessionid=832EE576C867695DB1B00600E0B8FCC7.d01t02?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Kim, S., Koo, I., Fang, A., and Zhang, X. (2011). Smith-Waterman peak alignment for comprehensive two-dimensional gas chromatography-mass spectrometry. *BMC Bioinformatics*, 12(1):235+. <http://www.biomedcentral.com/1471-2105/12/235>
- Koek, M. M., Kloet, F. M., Kleemann, R., Kooistra, T., Verheij, E. R., and Hankemeier, T. (2011). Semi-automated non-target processing in GCx GC-MS metabolomics analysis: applicability for biomedical studies. *Metabolomics*, 7(1):1–14. <http://www.springerlink.com/content/4036m33g1044v784/>
- Krupík, J., Májek, P., Gorovenko, R., Sandra, P., and Armstrong, D. W. (2011). On retentivity tuning by flow in the second column of different comprehensive two dimensional gas chromatographic configurations. *Journal of Chromatography A*, 1218(21):3186–3189. <http://www.sciencedirect.com/science/article/pii/S0021967311003906>
- Lei, Z., Huhman, D. V., and Sumner, L. W. (2011). Mass spectrometry strategies in metabolomics. *Journal of Biological Chemistry*, 286(29):25435–25442. <http://www.jbc.org/content/286/29/25435>
- Machado, M. E., Caramão, E. B., and Zini, C. A. (2011). Investigation of sulphur compounds in coal tar using monodimensional and comprehensive two-dimensional gas chromatography. *Journal of Chromatography A*, 1218(21):3200–3207. <http://www.sciencedirect.com/science/article/pii/S0021967310016651>
- Machado, M. E., Cappelli Fontanive, F., Oliveira, J. V., Caramão, E. B., and Alcaraz Zini, C. (2011). Identification of organic sulfur compounds in coal bitumen obtained by different extraction techniques using comprehensive two-dimensional gas chromatography coupled to time-of-flight mass spectrometric detection. *Analytical and Bioanalytical Chemistry*, 401(8):2433–2444. <http://www.springerlink.com/content/u515t656017u1838/>
- Mahé, L., Dutriez, T., Courtiade, M., Thiébaud, D., Dulot, H., and Bertoncini, F. (2011). Global approach for the selection of high temperature comprehensive two-dimensional gas chromatography experimental conditions and quantitative analysis in regards to sulfur-containing compounds in heavy petroleum cuts. *Journal of Chromatography A*, 1218(3):534–544. <http://www.sciencedirect.com/science/article/pii/S0021967310016535>
- McGregor, L. A., Gauchotte-Lindsay, C., Daéid, N. N., Thomas, R., Daly, P., and Kalin, R. M. (2011). Ultra resolution chemical fingerprinting of dense non-aqueous phase liquids from manufactured gas plants by reversed phase comprehensive two-dimensional gas chromatography. *Journal of Chromatography A*, 1218(29):4755–4763. <http://www.sciencedirect.com/science/article/pii/S002196731100687X>

- Mitrevski, B., Veleska, B., Engel, E., Wynne, P., Song, S. M., and Marriott, P. J. (2011). Chemical signature of ecstasy volatiles by comprehensive two-dimensional gas chromatography. *Forensic Science International*, 209(1-3):11–20. [http://www.fsijournal.org/article/S0379-0738\(10\)00509-8/abstract](http://www.fsijournal.org/article/S0379-0738(10)00509-8/abstract)
- Mol, H. G. J., van der Kamp, H., van der Weg, G., van der Lee, M., Punt, A., and de Rijk, T. C. 2011 Validation of automated Library-Based qualitative screening of pesticides by comprehensive Two-Dimensional gas Chromatography/Time-of-Flight mass spectrometry. *Journal of AOAC International*, pages 1722–1740. <http://www.ingentaconnect.com/content/aoac/jaoac/2011/00000094/00000006/art00006?token=00601f6d7f6b21ab3fc48227e442f2067217d6625445049762f5f7a42687627504541676249266d656c3b80788eff3a1>
- Muscalu, A. M., Reiner, E. J., Liss, S. N., Chen, T., Ladwig, G., and Morse, D. (2011). A routine accredited method for the analysis of polychlorinated biphenyls, organochlorine pesticides, chlorobenzenes and screening of other halogenated organics in soil, sediment and sludge by GCxGC- $\mu$ ECD. *Analytical and Bioanalytical Chemistry*, 401(8):2403–2413. <http://www.springerlink.com/content/c384gv81p74p4427/>
- Omais, B., Courtiade, M., Charon, N., Thiébaud, D., Quignard, A., and Hennion, M.-C. (2011). Investigating comprehensive two-dimensional gas chromatography conditions to optimize the separation of oxygenated compounds in a direct coal liquefaction middle distillate. *Journal of Chromatography A*, 1218(21):3233–3240. <http://www.sciencedirect.com/science/article/pii/S0021967310017413>
- Pani, O., Górecki, T., McNeish, C., Goldstein, A. H., Williams, B. J., Worton, D. R., Hering, S. V., and Kreisberg, N. M. (2011). Development of a new consumable-free thermal modulator for comprehensive two-dimensional gas chromatography. *Journal of Chromatography A*, 1218(20):3070–3079. <http://www.sciencedirect.com/science/article/pii/S0021967311003724>
- Parastar, H., Radovi, J. R., Jalali-Heravi, M., Diez, S., Bayona, J. M., and Tauler, R. (2011). Resolution and quantification of complex mixtures of polycyclic aromatic hydrocarbons in heavy fuel oil sample by means of GC xGC-TOFMS combined to multivariate curve resolution. *Anal. Chem.*, 83(24):9289–9297. <http://pubs.acs.org/doi/abs/10.1021/ac201799r>
- Patil, S. H., Banerjee, K., Utture, S. C., Fontana, A. R., Altamirano, J. C., Oulkar, D. P., Wagh, S. S., Dasgupta, S., Patil, S. B., Jadhav, M. R., Ugare, B. R., Adsule, P. G., and Deshmukh, M. B. (2011). Development and validation of a simple analytical method for the determination of 2,4,6-trichloroanisole in wine by GC-MS. *Food Chemistry*, 124(4):1734–1740. <http://www.sciencedirect.com/science/article/pii/S0308814610009817>
- Payeur, A. L., Meyers, P. A., and Sacks, R. D. (2011). Evaluation of on-line pyrolysis two-dimensional gas chromatography time-of-flight mass spectrometry (Py-GCxGC-ToFMS) on whole sediments from a mediterranean sapropel sequence. *Organic Geochemistry*, 42(10):1263–1270. <http://www.sciencedirect.com/science/article/pii/S014663801100204X>
- Pena-Abaurrea, M., Covaci, A., and Ramos, L. (2011). Comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry for the identification of organobrominated compounds in bluefin tuna. *Journal of Chromatography A*, 1218(39):6995–7002. <http://www.sciencedirect.com/science/article/pii/S0308814610009817>
- Petersen, M. A. and Grade, H. (2011). Analysis of steam assisted gravity drainage produced water using Two-Dimensional gas chromatography with Time-of-Flight mass spectrometry. *Ind. Eng. Chem. Res.*, 50(21):12217–12224. <http://pubs.acs.org/doi/abs/10.1021/ie200531h>
- Pierce, K. M. and Mohler, R. E. (2011). A review of chemometrics applied to comprehensive two-dimensional separations from 2008–2010. *Separation & Purification Reviews*, 41(2):143–168. <http://www.tandfonline.com/doi/abs/10.1080/15422119.2011.591868>
- Pleil, J. D. (2011). Non invasive biomedical analysis. breath networking session at PittCon 2011, atlanta, georgia. *Journal of Breath Research*, 5(2):029001+. <http://iopscience.iop.org/1752-7163/5/2/029001/>
- Ralston-Hooper, K., Jannasch, A., Adamec, J., and Sepúlveda, M. (2011a). The use of Two-Dimensional gas Chromatography-Time-of-Flight mass spectrometry (GCxGC-TOF-MS) for metabolomic analysis of polar metabolites metabolic profiling. volume 708 of *Methods in Molecular Biology*, chapter 12, pages 205–211. Humana Press, Totowa, NJ. <http://www.springerlink.com/content/j425150218371h65/#section=837872&page=1>
- Ralston-Hooper, K. J., Adamec, J., Jannasch, A., Mollenhauer, R., Ochoa-Acuña, H., and Sepúlveda, M. S. (2011b). Use of GC xGC/TOF-MS and LC/TOF-MS for metabolomic analysis of hyalella azteca chronically exposed to atrazine and its primary metabolite, desethylatrazine. *J. Appl. Toxicol.*, 31(5):399–410. <http://onlinelibrary.wiley.com/doi/10.1002/jat.1587/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Robinson, Adams, Boss, Heymann, H., Solomon, and Trengove (2011a). The relationship between sensory attributes and wine composition for australian cabernet sauvignon wines. *Australian Journal of Grape and Wine Research*, 17(3):327–340. <http://onlinelibrary.wiley.com/doi/10.1111/j.1755-0238.2011.00155.x/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Robinson, A. L., Boss, P. K., Heymann, H., Solomon, P. S., and Trengove, R. D. (2011b). Development of a sensitive non-targeted method for characterizing the wine volatile profile using headspace solid-phase microextraction comprehensive two-dimensional gas chromatography time-of-flight mass spectrometry. *Journal of Chromatography A*, 1218(3):504–517. <http://www.sciencedirect.com/science/article/pii/S0021967310015542>
- Robinson, A. L., Boss, P. K., Heymann, H., Solomon, P. S., and Trengove, R. D. (2011c). Influence of yeast strain, canopy management, and site on the volatile composition and sensory attributes of cabernet sauvignon wines from western australia. *J. Agric. Food Chem.*, 59(7):3273–3284. <http://pubs.acs.org/doi/abs/10.1021/jf104324d>



- Ruiz-Jimenez, J., Parshintsev, J., Laitinen, T., Hartonen, K., Riekkola, M.-L., Petaja, T., and Kulmala, M. (2011). Comprehensive two-dimensional gas chromatography, a valuable technique for screening and semiquantitation of different chemical compounds in ultrafine 30 nm and 50 nm aerosol particles. *J. Environ. Monit.*, 13(11):2994–3003. <http://pubs.rsc.org/en/Content/ArticleLanding/2011/EM/c1em10486a>
- Schnelle-Kreis, J., Orasche, J., Abbaszade, G., Schäfer, K., Harlos, D. P., Hansen, A. D. A., and Zimmermann, R. (2011). Application of direct thermal desorption gas chromatography time-of-flight mass spectrometry for determination of nonpolar organics in low-volume samples from ambient particulate matter and personal samplers. *Analytical and Bioanalytical Chemistry*, 401(10):3083–3094. <http://www.springerlink.com/content/0273024888vw6245/>
- Semard, G., Gouin, C., Bourdet, J., Bord, N., and Livadaris, V. (2011). Comparative study of differential flow and cryogenic modulators comprehensive two-dimensional gas chromatography systems for the detailed analysis of light cycle oil. *Journal of Chromatography A*, 1218(21):3146–3152. <http://www.sciencedirect.com/science/article/pii/S0021967310012045>
- Sfetsas, T., Michailof, C., Lappas, A., Li, Q., and Kneale, B. (2011). Qualitative and quantitative analysis of pyrolysis oil by gas chromatography with flame ionization detection and comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometry. *Journal of Chromatography A*, 1218(21):3317–3325. <http://www.sciencedirect.com/science/article/pii/S002196731001397X>
- Silva, R. S. F., Aguiar, H. G. M., Rangel, M. D., Azevedo, D. A., and Aquino Neto, F. R. (2011). Comprehensive two-dimensional gas chromatography with time of flight mass spectrometry applied to biomarker analysis of oils from colombia. *Fuel*, 90(8):2694–2699. <http://www.sciencedirect.com/science/article/pii/S001623611100233X>
- Skogerson, K., Wohlgemuth, G., Barupal, D., and Fiehn, O. (2011). The volatile compound BinBase mass spectral database. *BMC Bioinformatics*, 12(1):321+. <http://www.biomedcentral.com/1471-2105/12/321>
- Spikmans, V., Vaughan, G., Guo, C., and Fuller, S. (2011). Using sterol profiles for fingerprinting biodiesel and matching biodiesel spill samples to a source. *Environmental Forensics*, 12(1):19–24. <http://www.tandfonline.com/doi/abs/10.1080/15275922.2010.547548>
- Sun, Q., Gates, M. J., Lavin, E. H., Acree, T. E., and Sacks, G. L. (2011). Comparison of Odor-Active compounds in grapes and wines from vitis vinifera and Non-Foxy american grape species. *J. Agric. Food Chem.*, 59(19):10657–10664. <http://pubs.acs.org/doi/abs/10.1021/jf2026204>
- Tuan Anuar, S., Villegas, C., Mugo, S. M., and Curtis, J. M. (2011). The development of flow-through Bio-Catalyst microreactors from silica micro structured fibers for lipid transformations. *Lipids*, 46(6):545–555. <http://www.springerlink.com/content/2172267311648677/>
- van Stee, L. L. P. and Brinkman, U. A. (2011). Peak clustering in two-dimensional gas chromatography with mass spectrometric detection based on theoretical calculation of two-dimensional peak shapes: The 2DAid approach. *Journal of Chromatography A*, 1218(43):7878–7885. <http://www.sciencedirect.com/science/article/pii/S0021967311013094>
- van der Westhuizen, R., Potgieter, H., Prinsloo, N., de Villiers, A., and Sandra, P. (2011). Fractionation by liquid chromatography combined with comprehensive two-dimensional gas chromatography-mass spectrometry for analysis of cyclics in oligomerisation products of Fischer-Tropsch derived light alkenes. *Journal of Chromatography A*, 1218(21):3173–3179. <http://www.sciencedirect.com/science/article/pii/S0021967310013579>
- van der Westhuizen, R., Ajam, M., De Coning, P., Beens, J., de Villiers, A., and Sandra, P. (2011). Comprehensive two-dimensional gas chromatography for the analysis of synthetic and crude-derived jet fuels. *Journal of Chromatography A*, 1218(28):4478–4486. <http://www.sciencedirect.com/science/article/pii/S0021967311006376>
- Ventura, G. T., Hall, G. J., Nelson, R. K., Frysinger, G. S., Raghuraman, B., Pomerantz, A. E., Mullins, O. C., and Reddy, C. M. (2011). Analysis of petroleum compositional similarity using multiway principal components analysis (MPCA) with comprehensive two-dimensional gas chromatographic data. *Journal of Chromatography A*, 1218(18):2584–2592. <http://www.sciencedirect.com/science/article/pii/S0021967311003244>
- Vestner, J., Malherbe, S., Du Toit, M., Nieuwoudt, H. H., Mostafa, A., Górecki, T., Tredoux, A. G. J., and de Villiers, A. (2011). Investigation of the volatile composition of pinotage wines fermented with different malolactic starter cultures using comprehensive Two-Dimensional gas chromatography coupled to Time-of-Flight mass spectrometry (GCxGC-TOF-MS). *J. Agric. Food Chem.*, 59(24):12732–12744. <http://pubs.acs.org/doi/abs/10.1021/jf2028208>
- Waldhier, M. C., Almstetter, M. F., Nürnberger, N., Gruber, M. A., Dettmer, K., and Oefner, P. J. (2011). Improved enantiomer resolution and quantification of free d-amino acids in serum and urine by comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry. *Journal of Chromatography A*, 1218(28):4537–4544. <http://www.sciencedirect.com/science/article/pii/S0021967311006674>
- Wang, H., Weng, N., Zhang, S., Zhu, G., and Wei, C. (2011). Comparison of geochemical parameters derived from comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometry and conventional gas chromatography-mass spectrometry. *SCIENCE CHINA Earth Sciences*, 54(12):1892–1901. <http://www.springerlink.com/content/g4w5257030241107/>
- Weldegergis, B. T., Villiers, A., McNeish, C., Seethapathy, S., Mostafa, A., Górecki, T., and Crouch, A. M. (2011). Characterisation of volatile components of pinotage wines using comprehensive two-dimensional gas chromatography coupled to time-of-flight mass spectrometry (GCxGC-TOFMS). *Food Chemistry*, 129(1):188–199. <http://www.sciencedirect.com/science/article/pii/S0308814610015852>
- Wong, J. W., Zhang, K., Hayward, D. G., and Kai-Meng, C. (2011). Multiresidue pesticide analysis by capillary gas Chromatography-Mass spectrometry mass spectrometry in food safety. volume 747 of *Methods in Molecular Biology*, chapter 6, pages 131–172. Humana Press, Totowa, NJ. <http://www.springerlink.com/content/xt155117867001v8/#section=902155&page=1>

- Zapadlo, M., Krupik, J., Kovalczuk, T., Májek, P., pánik, I., Armstrong, D. W., and Sandra, P. (2011). Enhanced comprehensive two-dimensional gas chromatographic resolution of polychlorinated biphenyls on a non-polar polysiloxane and an ionic liquid column series. *Journal of Chromatography A*, 1218(5):746–751. <http://www.sciencedirect.com/science/article/pii/S0021967310016894>
- Zeng, Z.-D., Hugel, H. M., and Marriott, P. J. (2011). Chemometrics in comprehensive multidimensional separations. *Analytical and Bioanalytical Chemistry*, 401(8):2373–2386. <http://www.springerlink.com/content/g66855150u246442/>
- Zhang, S., Zhang, B., Zhu, G., Wang, H., and Li, Z. (2011). Geochemical evidence for coal-derived hydrocarbons and their charge history in the dabei gas field, kuqa thrust belt, tarim basin, NW china. *Marine and Petroleum Geology*, 28(7):1364–1375. <http://www.sciencedirect.com/science/article/pii/S0264817211000717>
- Zhao, Y., Zhang, J., Wang, B., Kim, S. H., Fang, A., Bogdanov, B., Zhou, Z., McClain, C., and Zhang, X. (2011). A method of calculating the second dimension retention index in comprehensive two-dimensional gas chromatography time-of-flight mass spectrometry. *Journal of Chromatography A*, 1218(18):2577–2583. <http://www.sciencedirect.com/science/article/pii/S0021967311003177>
- Zrostlikova, J., Binkley, J., Wojtowicz, P., and Adam, T. (2011). Application of comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometry in human metabolomic studies. *Journal of Analytical Chemistry*, 66(13):1333–1340. <http://www.springerlink.com/content/n382182u50366t2k/>

## 2010

---

- Araújo, R. C., Pasa, V. M., Marriott, P. J., and Cardeal, Z. L. (2010). Analysis of volatile organic compounds in polyurethane coatings based on eucalyptus sp. bio-oil pitch using comprehensive two-dimensional gas chromatography (GCxGC). *Journal of Analytical and Applied Pyrolysis*, 88(1):91–97. <http://www.sciencedirect.com/science/article/pii/S0165237010000331>
- Asiago, V. M., Alvarado, L. Z., Shanaiah, N., Gowda, G. A. N., Owusu-Sarfo, K., Ballas, R. A., and Raftery, D. (2010). Early detection of recurrent breast cancer using metabolite profiling. *Cancer Research*, 70(21):8309–8318. <http://cancerres.aacrjournals.org/content/70/21/8309>
- Ávila, B. M. F., Aguiar, A., Gomes, A. O., and Azevedo, D. A. (2010). Characterization of extra heavy gas oil biomarkers using comprehensive two-dimensional gas chromatography coupled to time-of-flight mass spectrometry. *Organic Geochemistry*, 41(9):863–866. <http://www.sciencedirect.com/science/article/pii/S0146638010000768>
- Baharum, S. N., Bunawan, H., Abd, M., Mustapha, W. A. W., and Noor, N. M. (2010). Analysis of the chemical composition of the essential oil of polygonum minus huds. using Two-Dimensional gas Chromatography-Time-of-Flight mass spectrometry (GC-TOF MS). *Molecules*, 15(10):7006–7015. <http://www.mdpi.com/1420-3049/15/10/7006>
- Cajka, T., Riddellova, K., Klimankova, E., Cerna, M., Pudil, F., and Hajslova, J. (2010). Traceability of olive oil based on volatiles pattern and multivariate analysis. *Food Chemistry*, 121(1):282–289. <http://www.sciencedirect.com/science/article/pii/S0308814609014101>
- Cooper, S. J., Finney, G. L., Brown, S. L., Nelson, S. K., Hesselberth, J., MacCoss, M. J., and Fields, S. (2010). High-throughput profiling of amino acids in strains of the *saccharomyces cerevisiae* deletion collection. *Genome Research*, 20(9):1288–1296. <http://genome.cshlp.org/content/20/9/1288>
- Costa, R., Dugo, P., Santi, L., Dugo, G., and Mondello, L. (2010). Advances of modern gas chromatography and hyphenated techniques for analysis of plant extracts. *Current Organic Chemistry*, pages 1752–1768. <http://www.ingentaconnect.com/content/ben/coc/2010/00000014/00000016/art00007?token=00561a48545c217f177e2a46762c6b412148704670534979457a673f7b2f267738703375686f495b0f702c>
- Doebbe, A., Keck, M., La Russa, M., Mussgnug, J. H., Hankamer, B., Tekçe, E., Niehaus, K., and Kruse, O. (2010). The interplay of proton, electron, and metabolite supply for photosynthetic h<sub>2</sub> production in *chlamydomonas reinhardtii*. *Journal of Biological Chemistry*, 285(39):30247–30260. <http://www.jbc.org/content/285/39/30247>
- Dutriez, T., Courtiade, M., Thiébaud, D., Dulot, H., Bertoncini, F., and Hennion, M.-C. (2010). Extended characterization of a vacuum gas oil by offline LC-high-temperature comprehensive two-dimensional gas chromatography. *J. Sep. Science*, 33(12):1787–1796. <http://onlinelibrary.wiley.com/doi/10.1002/jssc.201000102/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Gao, X., Williams, S. J., Woodman, O. L., and Marriott, P. J. (2010). Comprehensive two-dimensional gas chromatography, retention indices and time-of-flight mass spectra of flavonoids and chalcones. *Journal of Chromatography A*, 1217(52):8317–8326. <http://www.sciencedirect.com/science/article/pii/S0021967310014883>
- Guthery, B., Bassindale, T., Bassindale, A., Pillinger, C. T., and Morgan, G. H. (2010). Qualitative drug analysis of hair extracts by comprehensive two-dimensional gas chromatography/time-of-flight mass spectrometry. *Journal of Chromatography A*, 1217(26):4402–4410. <http://www.sciencedirect.com/science/article/pii/S0021967310004887>
- Hayward, D. G., Wong, J. W., and Zhang, K. 2010 Multiresidue pesticide analysis in ginseng and spinach by nontargeted and targeted screening procedures. *Journal of AOAC International*, pages 1741–1751. <http://www.ingentaconnect.com/content/aoac/jaoac/2011/00000094/00000006/art00007?token=005c1f99bcb29a8a5ce61166720297d76344470234a457a246a4d572b6d3f6a4b4b6e6e42576b642738f784c07a0>
- Hayward, D. G., Pisano, T. S., Wong, J. W., and Scudder, R. J. (2010). Multiresidue method for pesticides and persistent organic pollutants (POPs) in milk and cream using comprehensive Two-Dimensional capillary gas Chromatography-Time-of-Flight mass spectrometry. *J. Agric. Food Chem.*, 58(9):5248–5256. <http://pubs.acs.org/doi/abs/10.1021/jf100021p>
- Hilton, D. C., Jones, R. S., and Sjödin, A. (2010). A method for rapid, non-targeted screening for environmental contaminants in household dust. *Journal of Chromatography A*, 1217(44):6851–6856. <http://www.sciencedirect.com/science/article/pii/S0021967310011118>

- Indrasti, D., Che Man, Y. B., Mustafa, S., and Hashim, D. M. (2010). Lard detection based on fatty acids profile using comprehensive gas chromatography hyphenated with time-of-flight mass spectrometry. *Food Chemistry*, 122(4):1273–1277. <http://www.sciencedirect.com/science/article/pii/S0308814610003754>
- Indrasti, D., Che Man, Y. B., Chin, S. T., Mustafa, S., Mat Hashim, D., and Abdul Manaf, M. (2010). Regiospecific analysis of mono- and diglycerides in glycerolysis products by GC x GC-TOF-MS. *Journal of the American Oil Chemists' Society*, 87(11):1255–1262. <http://www.springerlink.com/content/x4127j4746562844/>
- Kind, T. and Fiehn, O. (2010). Advances in structure elucidation of small molecules using mass spectrometry. *Bioanalytical Reviews*, 2(1):23–60. <http://www.springerlink.com/content/c1p83306688q7656/>
- Klein, M. S., Almstetter, M. F., Schlamberger, G., Nürnberger, N., Dettmer, K., Oefner, P. J., Meyer, H. H. D., Wiedemann, S., and Gronwald, W. (2010). Nuclear magnetic resonance and mass spectrometry-based milk metabolomics in dairy cows during early and late lactation. *Journal of Dairy Science*, 93(4):1539–1550. [http://www.journalofdairyscience.org/article/S0022-0302\(10\)00127-X/abstract](http://www.journalofdairyscience.org/article/S0022-0302(10)00127-X/abstract)
- Kohl, A., Cochran, J., and Cropek, D. M. (2010). Characterization of military fog oil by comprehensive two-dimensional gas chromatography. *Journal of Chromatography A*, 1217(4):550–557. <http://www.sciencedirect.com/science/article/pii/S0021967309017336>
- Kouremenos, K. A., Harynuk, J. J., Winniford, W. L., Morrison, P. D., and Marriott, P. J. (2010a). One-pot microwave derivatization of target compounds relevant to metabolomics with comprehensive two-dimensional gas chromatography. *Journal of Chromatography B*, 878(21):1761–1770. <http://www.sciencedirect.com/science/article/pii/S1570023210002850>
- Kouremenos, K. A., Pitt, J., and Marriott, P. J. (2010b). Metabolic profiling of infant urine using comprehensive two-dimensional gas chromatography: Application to the diagnosis of organic acidurias and biomarker discovery. *Journal of Chromatography A*, 1217(1):104–111. <http://www.sciencedirect.com/science/article/pii/S0021967309015313>
- Laitinen, T., Martín, S. H., Parshintsev, J., Hyötyläinen, T., Hartonen, K., Riekkola, M.-L., Kulmala, M., and Pavón, J. L. (2010). Determination of organic compounds from wood combustion aerosol nanoparticles by different gas chromatographic systems and by aerosol mass spectrometry. *Journal of Chromatography A*, 1217(1):151–159. <http://www.sciencedirect.com/science/article/pii/S0021967309016896>
- Lee, J. and Lane, D. A. (2010). Formation of oxidized products from the reaction of gaseous phenanthrene with the OH radical in a reaction chamber. *Atmospheric Environment*, 44(20):2469–2477. <http://www.sciencedirect.com/science/article/pii/S1352231010002037>
- Li, Y., Pang, T., Guo, Z., Li, Y., Wang, X., Deng, J., Zhong, K., Lu, X., and Xu, G. (2010). Accelerated solvent extraction for GC-based tobacco fingerprinting and its comparison with simultaneous distillation and extraction. *Talanta*, 81(1-2):650–656. <http://www.sciencedirect.com/science/article/pii/S0039914010000135>
- Libardoni, M., Fix, C., Waite, J. H., and Sacks, R. (2010). Design and performance evaluation of a two-stage resistively-heated thermal modulator for GC [times] GC. *Anal. Methods*, 2(7):936–943. <http://pubs.rsc.org/en/Content/ArticleLanding/2010/AY/c0ay00090f>
- Matamoros, V., Jover, E., and Bayona, J. M. (2010). Occurrence and fate of benzothiazoles and benzotriazoles in constructed wetlands. *Water Science & Technology*, 61(1):191+. <http://www.iwaponline.com/wst/06101/wst061010191.htm>
- Mateus, E., Barata, R. C., Zrostlíková, J., Gomes da Silva, M. D. R., and Paiva, M. R. (2010a). Characterization of the volatile fraction emitted by pinus spp. by one- and two-dimensional chromatographic techniques with mass spectrometric detection. *Journal of Chromatography A*, 1217(11):1845–1855. <http://www.sciencedirect.com/science/article/pii/S0021967310000828>
- Mateus, E. P., Zrostlíková, J., Gomes da Silva, M. D. R., Ribeiro, A. B., and Marriott, P. (2010b). Electrokinetic removal of creosote from treated timber waste: a comprehensive gas chromatographic view. *Journal of Applied Electrochemistry*, 40(6):1183–1193. <http://www.springerlink.com/content/j670278t256401p7/>
- Mieth, M., Schubert, J. K., Großger, T., Sabel, B., Kischkel, S., Fuchs, P., Hein, D., Zimmermann, R., and Miekisch, W. (2010). Automated needle trap Heart-Cut GC/MS and needle trap comprehensive Two-Dimensional GC/TOF-MS for breath gas analysis in the clinical environment. *Anal. Chem.*, 82(6):2541–2551. <http://pubs.acs.org/doi/abs/10.1021/ac100061k>
- Mitreviski, B. S., Wilairat, P., and Marriott, P. J. (2010). Evaluation of world Anti-Doping agency criteria for anabolic agent analysis by using comprehensive two-dimensional gas chromatography mass spectrometry. *Analytical and Bioanalytical Chemistry*, 396(7):2503–2511. <http://www.springerlink.com/content/e03718v162118087/>
- Natsch, A., Kuhn, F., and Tiercy, J.-M. (2010). Lack of evidence for HLA-linked patterns of odorous carboxylic acids released from glutamine conjugates secreted in the human axilla. *Journal of Chemical Ecology*, 36(8):837–846. <http://www.springerlink.com/content/etw48324mg622028/>
- Okubo, Y., Yang, S., Chistoserdova, L., and Lidstrom, M. E. (2010). Alternative route for glyoxylate consumption during growth on Two-Carbon compounds by methylotrophic bacterium *Methylobacterium extorquens* AM1. *Journal of Bacteriology*, 192(7):1813–1823. <http://jb.asm.org/content/192/7/1813>
- Ozel, M. Z., Gogus, F., Yagci, S., Hamilton, J. F., and Lewis, A. C. (2010). Determination of volatile nitrosamines in various meat products using comprehensive gas chromatography-nitrogen chemiluminescence detection. *Food and Chemical Toxicology*, 48(11):3268–3273. <http://www.sciencedirect.com/science/article/pii/S0278691510005570>
- Özel, M. Z., Ward, M. W., Hamilton, J. F., Lewis, A. C., Raventós-Duran, T., and Harrison, R. M. (2010). Analysis of organic nitrogen compounds in urban aerosol samples using GCxGC-TOF/MS. *Aerosol Science and Technology*, 44(2):109–116. <http://www.tandfonline.com/doi/abs/10.1080/02786820903410105>

- Peacock, E. E., Arey, J. S., DeMello, J. A., McNichol, A. P., Nelson, R. K., and Reddy, C. M. (2010). Molecular and isotopic analysis of motor oil from a Biodiesel-Driven vehicle. *Energy Fuels*, 24(2):1037–1042. <http://pubs.acs.org/doi/abs/10.1021/ef9011388>
- Perestrelo, R., Petronilho, S., Câmara, J. S., and Rocha, S. M. (2010). Comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometry combined with solid phase microextraction as a powerful tool for quantification of ethyl carbamate in fortified wines. the case study of madeira wine. *Journal of Chromatography A*, 1217(20):3441–3445. <http://www.sciencedirect.com/science/article/pii/S0021967310003912>
- Risticvic, S., Chen, Y., Kudlejova, L., Vatinno, R., Baltensperger, B., Stuff, J. R., Hein, D., and Pawliszyn, J. (2010). Protocol for the development of automated high-throughput SPME-GC methods for the analysis of volatile and semivolatile constituents in wine samples. *Nature Protocols*, 5(1):162–176. [http://pubget.com/paper/20057386/Protocol\\_for\\_the\\_development\\_of\\_automated\\_high\\_throughput\\_SPME\\_GC\\_methods\\_for\\_the\\_analysis\\_of\\_volatile\\_and\\_semivolatile\\_constituents\\_in\\_wine\\_samples](http://pubget.com/paper/20057386/Protocol_for_the_development_of_automated_high_throughput_SPME_GC_methods_for_the_analysis_of_volatile_and_semivolatile_constituents_in_wine_samples)
- Rosse, A. D. and Harynyuk, J. J. (2010). Methodology for the derivatization and quantification of dialkyl phosphate esters in petroleum samples. *Anal. Methods*, 2(8):1176–1179. <http://pubs.rsc.org/en/Content/ArticleLanding/2010/AY/c0ay00235f>
- Scheiner, J. J., Sacks, G. L., Pan, B., Ennahli, S., Tarlton, L., Wise, A., Lerch, S. D., and Vanden Heuvel, J. E. (2010). Impact of severity and timing of basal leaf removal on 3-Isobutyl-2-Methoxypyrazine concentrations in red winegrapes. *American Journal of Enology and Viticulture*, 61(3):358–364. <http://www.ajevonline.org/content/61/3/358.abstract>
- Schmarr, H.-G. and Bernhardt, J. (2010). Profiling analysis of volatile compounds from fruits using comprehensive two-dimensional gas chromatography and image processing techniques. *Journal of Chromatography A*, 1217(4):565–574. <http://www.sciencedirect.com/science/article/pii/S0021967309017427>
- Semard, G., Peulon-Agasse, V., Bruchet, A., Bouillon, J.-P., and Cardinaël, P. (2010). Convex hull: A new method to determine the separation space used and to optimize operating conditions for comprehensive two-dimensional gas chromatography. *Journal of Chromatography A*, 1217(33):5449–5454. <http://www.sciencedirect.com/science/article/pii/S0021967310008356>
- Shepherd, A. G., van Mispelaar, V., Nowlin, J., Genuit, W., and Grutters, M. (2010). Analysis of naphthenic acids and derivatization agents using Two-Dimensional gas chromatography and mass spectrometry: Impact on flow assurance prediction†. *Energy Fuels*, 24(4):2300–2311. <http://pubs.acs.org/doi/abs/10.1021/ef900949m>
- Snyder, L. R., Hoggard, J. C., Montine, T. J., and Synovec, R. E. (2010). Development and application of a comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometry method for the analysis of L-β-methylamino-alanine in human tissue. *Journal of Chromatography A*, 1217(27):4639–4647. <http://www.sciencedirect.com/science/article/pii/S0021967310005832>
- Stanimirova, I., Üstün, B., Cajka, T., Riddelova, K., Hajslova, J., Buydens, L. M. C., and Walczak, B. (2010). Tracing the geographical origin of honeys based on volatile compounds profiles assessment using pattern recognition techniques. *Food Chemistry*, 118(1):171–176. <http://www.sciencedirect.com/science/article/pii/S0308814609005457>
- Théron, L., Tournayre, P., Kondjoyan, N., Abouelkaram, S., Santé-Lhoutellier, V., and Berdagué, J.-L. (2010). Analysis of the volatile profile and identification of odour-active compounds in bayonne ham. *Meat Science*, 85(3):453–460. <http://www.sciencedirect.com/science/article/pii/S0039914010000135>
- Tran, T. C., Logan, G. A., Grosjean, E., Ryan, D., and Marriott, P. J. (2010). Use of comprehensive two-dimensional gas chromatography/time-of-flight mass spectrometry for the characterization of biodegradation and unresolved complex mixtures in petroleum. *Geochimica et Cosmochimica Acta*, 74(22):6468–6484. <http://www.sciencedirect.com/science/article/pii/S0016703710004734>
- van der Westhuizen, R., Crous, R., de Villiers, A., and Sandra, P. (2010). Comprehensive two-dimensional gas chromatography for the analysis of Fischer-Tropsch oil products. *Journal of Chromatography A*, 1217(52):8334–8339. <http://www.sciencedirect.com/science/article/pii/S0021967310014949>
- Velagapudi, V. R., Hezaveh, R., Reigstad, C. S., Gopalacharyulu, P., Yetukuri, L., Islam, S., Felin, J., Perkins, R., Borén, J., Orei, M., and Bäckhed, F. (2010). The gut microbiota modulates host energy and lipid metabolism in mice. *Journal of Lipid Research*, 51(5):1101–1112. <http://www.jlr.org/content/51/5/1101>
- Villegas, C., Zhao, Y., and Curtis, J. M. (2010). Two methods for the separation of monounsaturated octadecenoic acid isomers. *Journal of Chromatography A*, 1217(5):775–784. <http://www.sciencedirect.com/science/article/pii/S002196730901797X>
- Vivó-Truyols, G. and Janssen, H.-G. (2010). Probability of failure of the watershed algorithm for peak detection in comprehensive two-dimensional chromatography. *Journal of Chromatography A*, 1217(8):1375–1385. <http://www.sciencedirect.com/science/article/pii/S002196730901913X>
- von Muhlen, C., de Oliveira, E. C., Zini, C. A., Caramalfo, E. B., and Marriott, P. J. (2010). Characterization of Nitrogen-Containing compounds in heavy gas oil petroleum fractions using comprehensive Two-Dimensional gas chromatography coupled to Time-of-Flight mass spectrometry. *Energy Fuels*, 24(6):3572–3580. <http://pubs.acs.org/doi/abs/10.1021/ef1002364>
- Wahl, J. H. and Colburn, H. A. (2010). Extraction of chemical impurities for forensic investigations: A case study for indoor releases of a sarin surrogate. *Building and Environment*, 45(5):1339–1345. <http://www.sciencedirect.com/science/article/pii/S0360132309003205>
- Wang, B., Fang, A., Heim, J., Bogdanov, B., Pugh, S., Libardoni, M., and Zhang, X. (2010a). DISCO: Distance and spectrum correlation optimization alignment for Two-Dimensional gas chromatography Time-of-Flight mass Spectrometry-Based metabolomics. *Anal. Chem.*, 82(12):5069–5081. <http://pubs.acs.org/doi/abs/10.1021/ac100064b>



- Wang, H., Weng, N., Zhang, S., Zhu, G., Chen, J., and Wei, C. (2010b). Identification of petroleum aromatic fraction by comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometry. *Chinese Science Bulletin*, 55(19):2039–2045. <http://www.springerlink.com/content/p71717m183306g21/>
- Watson, J. S., Sephton, M. A., and Gilmour, I. (2010). Thermochemolysis of the murchison meteorite: identification of oxygen bound and occluded units in the organic macromolecule. *International Journal of Astrobiology*, 9(Special Issue 04):201–208. <http://journals.cambridge.org/action/displayAbstract?fromPage=online&aid=7909953>
- Wojtowicz, P., Zrostliková, J., Kovalczuk, T., Schrek, J., and Adam, T. (2010). Evaluation of comprehensive two-dimensional gas chromatography coupled to time-of-flight mass spectrometry for the diagnosis of inherited metabolic disorders using an automated data processing strategy. *Journal of Chromatography A*, 1217(51):8054–8061. <http://www.sciencedirect.com/science/article/pii/S0021967310013178>
- Wulf, V., Wienand, N., Wirtz, M., Kling, H.-W., Gäb, S., and Schmitz, O. J. (2010). Analysis of special surfactants by comprehensive two-dimensional gas chromatography coupled to time-of-flight mass spectrometry. *Journal of Chromatography A*, 1217(5):749–754. <http://www.sciencedirect.com/science/article/pii/S0021967309017804>

## 2009

---

- Adam, F., Bertoncini, F., Dartiguelongue, C., Marchand, K., Thiébaud, D., and Hennion, M.-C. (2009). Comprehensive two-dimensional gas chromatography for basic and neutral nitrogen speciation in middle distillates. *Fuel*, 88(5):938–946. <http://www.sciencedirect.com/science/article/pii/S0016236108004900>
- Almstetter, M. F., Appel, I. J., Gruber, M. A., Lottaz, C., Timischl, B., Spang, R., Dettmer, K., and Oefner, P. J. (2009). Integrative normalization and comparative analysis for metabolic fingerprinting by comprehensive Two-Dimensional gas Chromatography-Time-of-Flight mass spectrometry. *Anal. Chem.*, 81(14):5731–5739. <http://pubs.acs.org/doi/abs/10.1021/ac900528b>
- Amorim, L. C., Dimandja, J.-M., and Cardeal, Z. (2009). Analysis of hydroxylated polycyclic aromatic hydrocarbons in urine using comprehensive two-dimensional gas chromatography with a flame ionization detector. *Journal of Chromatography A*, 1216(14):2900–2904. <http://www.sciencedirect.com/science/article/pii/S0021967308019663>
- Begnaud, F., Debonneville, C., Probst, J.-P., Chaintreau, A., Morrison, P. D., Adcock, J. L., and Marriott, P. J. (2009). Effects of variation in modulator temperature during cryogenic modulation in comprehensive two-dimensional gas chromatography. *J. Sep. Science*, 32(18):3144–3151. <http://onlinelibrary.wiley.com/doi/10.1002/jssc.200900290/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Bertoncini, F., Marion, M. C., Brodusch, N., and Esnault, S. (2009). Unravelling molecular composition of products from cobalt catalysed Fischer-Tropsch reaction by comprehensive gas chromatography: Methodology and application. *Oil & Gas Science and Technology - Revue de l'IFP*, 64(1):79–90. [http://ogst.ifpenergiesnouvelles.fr/index.php?option=com\\_article&access=doi&doi=10.2516/ogst/2008054&Itemid=129](http://ogst.ifpenergiesnouvelles.fr/index.php?option=com_article&access=doi&doi=10.2516/ogst/2008054&Itemid=129)
- Bester, K. (2009). Analysis of musk fragrances in environmental samples. *Journal of Chromatography A*, 1216(3):470–480. <http://www.sciencedirect.com/science/article/pii/S0021967308014891>
- Bouvier, C., Reumkens, N., and Buijs, W. (2009). Separation of diisopropylnaphthalene isomers. *Journal of Chromatography A*, 1216(36):6410–6416. <http://www.sciencedirect.com/science/article/pii/S0021967309010243>
- Breme, K., Tournayre, P., Fernandez, X., Meierhenrich, U. J., Brevard, H., Joulain, D., and Berdaguel, J. L. (2009a). Characterization of volatile compounds of indian cress absolute by GC-Olfactometry/VIDEO-sniff and comprehensive Two-Dimensional gas chromatography. *J. Agric. Food Chem.*, 58(1):473–480. <http://pubs.acs.org/doi/abs/10.1021/jf902946v>
- Breme, K., Tournayre, P., Fernandez, X., Meierhenrich, U. J., Brevard, H., Joulain, D., and Berdaguel, J. L. (2009b). Identification of odor impact compounds of tagetes minuta L. essential oil: Comparison of two GC-olfactometry methods. *J. Agric. Food Chem.*, 57(18):8572–8580. <http://pubs.acs.org/doi/abs/10.1021/jf9016509>
- Cardeal, Z. L. and Marriott, P. J. (2009). Comprehensive two-dimensional gas chromatography-mass spectrometry analysis and comparison of volatile organic compounds in brazilian cachaça and selected spirits. *Food Chemistry*, 112(3):747–755. <http://www.sciencedirect.com/science/article/pii/S0308814608007528>
- Chin, S.-T., Che Man, Y. B., Tan, C. P., and Hashim, D. M. (2009). Rapid profiling of Animal-Derived fatty acids using fast GCxGC coupled to Time-of-Flight mass spectrometry. *Journal of the American Oil Chemists' Society*, 86(10):949–958. <http://www.springerlink.com/content/4h0l56j856406694/>
- de Souza, P. P., Cardeal, Z., Augusti, R., Morrison, P., and Marriott, P. J. (2009). Determination of volatile compounds in brazilian distilled cachaça by using comprehensive two-dimensional gas chromatography and effects of production pathways. *Journal of Chromatography A*, 1216(14):2881–2890. <http://www.sciencedirect.com/science/article/pii/S0021967308018311>
- Eganhouse, R. P., Pontolillo, J., Gaines, R. B., Frysinger, G. S., Gabriel, F. L. P., Kohler, H.-P. E., Giger, W., and Barber, L. B. (2009). Isomer-Specific determination of 4-Nonylphenols using comprehensive Two-Dimensional gas Chromatography/Time-of-Flight mass spectrometry. *Environ. Sci. Technol.*, 43(24):9306–9313. <http://pubs.acs.org/doi/abs/10.1021/es902622r>
- Freitas, L. S., Von Mühlen, C., Bortoluzzi, J. H., Zini, C. A., Fortuny, M., Dariva, C., Coutinho, R. C. C., Santos, A. F., and Caramão, E. B. (2009). Analysis of organic compounds of water-in-crude oil emulsions separated by microwave heating using comprehensive two-dimensional gas chromatography and time-of-flight mass spectrometry. *Journal of Chromatography A*, 1216(14):2860–2865. <http://www.sciencedirect.com/science/article/pii/S0021967308015446>

- Gaquerel, E., Weinhold, A., and Baldwin, I. T. (2009). Molecular interactions between the specialist herbivore *Manduca sexta* (Lepidoptera, Sphingidae) and its natural host *Nicotiana attenuata*. VIII. An unbiased GCxGC-ToFMS analysis of the plant's elicited volatile emissions. *Plant Physiology*, 149(3):1408–1423. <http://www.plantphysiol.org/content/149/3/1408>
- Guthery, B., Bassindale, A., Pillinger, C. T., and Morgan, G. H. (2009). The detection of various opiates and benzodiazepines by comprehensive two-dimensional gas chromatography/time-of-flight mass spectrometry. *Rapid Commun. Mass Spectrom.*, 23(3):340–348. <http://onlinelibrary.wiley.com/doi/10.1002/rcm.3883/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Hoggard, J. C., Siegler, W. C., and Synovec, R. E. (2009). Toward automated peak resolution in complete GCxGC-TOFMS chromatograms by PARAFAC. *J. Chemometrics*, 23(7-8):421–431. <http://onlinelibrary.wiley.com/doi/10.1002/cem.1239/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Hoggard, J. C., Wahl, J. H., Synovec, R. E., Mong, G. M., and Fraga, C. G. (2009). Impurity profiling of a chemical weapon precursor for possible forensic signatures by comprehensive Two-Dimensional gas Chromatography/Mass spectrometry and chemometrics. *Anal. Chem.*, 82(2):689–698. <http://pubs.acs.org/doi/abs/10.1021/ac902247x>
- Hoh, E., Lehotay, S. J., Mastovska, K., Ngo, H. L., Vetter, W., Pangallo, K. C., and Reddy, C. M. (2009). Capabilities of direct sample Introduction-Comprehensive Two-Dimensional gas Chromatography-Time-of-Flight mass spectrometry to analyze organic chemicals of interest in fish oils. *Environ. Sci. Technol.*, 43(9):3240–3247. <http://pubs.acs.org/doi/abs/10.1021/es803486x>
- Humston, E. M., Hoggard, J. C., and Synovec, R. E. (2009). Utilizing the third order advantage with isotope dilution mass spectrometry. *Anal. Chem.*, 82(1):41–43. <http://pubs.acs.org/doi/abs/10.1021/ac902184b>
- Janssen, H.-G., Steenbergen, H., and de Koning, S. (2009). The role of comprehensive chromatography in the characterization of edible oils and fats. *Eur. J. Lipid Sci. Technol.*, 111(12):1171–1184. <http://onlinelibrary.wiley.com/doi/10.1002/ejlt.200900074/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Jover, E., Matamoros, V., and Bayona, J. M. (2009). Characterization of benzothiazoles, benzotriazoles and benzosulfonamides in aqueous matrixes by solid-phase extraction followed by comprehensive two-dimensional gas chromatography coupled to time-of-flight mass spectrometry. *Journal of Chromatography A*, 1216(18):4013–4019. <http://www.sciencedirect.com/science/article/pii/S002196730900260X>
- Kalinová, B., Podskalská, H., Rika, J., and Hoskovec, M. (2009). Irresistible bouquet of death-how are burying beetles (Coleoptera: Silphidae: Nicrophorus) attracted by carcasses. *Naturwissenschaften*, 96(8):889–899. <http://www.springerlink.com/content/r13g600624g3w88n/>
- Kempa, S., Hummel, J., Schwemmer, T., Pietzke, M., Strehmel, N., Wienkoop, S., Kopka, J., and Weckwerth, W. (2009). An automated GCxGC-TOF-MS protocol for batch-wise extraction and alignment of mass isotopomer matrixes from differential <sup>13</sup>C-labelling experiments: a case study for photoautotrophic-mixotrophic grown *Chlamydomonas reinhardtii* cells. *J. Basic Microbiol.*, 49(1):82–91. <http://onlinelibrary.wiley.com/doi/10.1002/jobm.200800337/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Koning, S., Janssen, H.-G., and Brinkman, U. A. (2009). Modern methods of sample preparation for GC analysis. *Chromatographia*, 69(0):33–78. <http://www.springerlink.com/content/637w27203k163377/>
- Lojzova, L., Riddellova, K., Hajslova, J., Zrostlikova, J., Schurek, J., and Cajka, T. (2009). Alternative GC-MS approaches in the analysis of substituted pyrazines and other volatile aromatic compounds formed during Maillard reaction in potato chips. *Analytica Chimica Acta*, 641(1-2):101–109. <http://www.sciencedirect.com/science/article/pii/S0003267009004437>
- Ma, C., Wang, H., Lu, X., Wang, H., Xu, G., and Liu, B. (2009). Terpenoid metabolic profiling analysis of transgenic *Artemisia annua* L. by comprehensive two-dimensional gas chromatography time-of-flight mass spectrometry. *Metabolomics*, 5(4):497–506. <http://www.springerlink.com/content/17g5732510702277/>
- Mao, D., Lookman, R., Weghe, H. V., Weltens, R., Vanermen, G., Brucker, N. D., and Diels, L. (2009). Estimation of ecotoxicity of petroleum hydrocarbon mixtures in soil based on HPLC-GC/MS analysis. *Chemosphere*, 77(11):1508–1513. <http://www.sciencedirect.com/science/article/pii/S0045653509011837>
- Mao, D., Lookman, R., Van De Weghe, H., Vanermen, G., De Brucker, N., and Diels, L. (2009a). Aqueous solubility calculation for petroleum mixtures in soil using comprehensive two-dimensional gas chromatography analysis data. *Journal of Chromatography A*, 1216(14):2873–2880. <http://www.sciencedirect.com/science/article/pii/S0021967308014398>
- Mao, D., Weghe, Lookman, R., Vanermen, G., Brucker, N. D., and Diels, L. (2009b). Resolving the unresolved complex mixture in motor oils using high-performance liquid chromatography followed by comprehensive two-dimensional gas chromatography. *Fuel*, 88(2):312–318. <http://www.sciencedirect.com/science/article/pii/S0016236108003256>
- Mastovska, K., Dorweiler, K. J., Lehotay, S. J., Wegscheid, J. S., and Szpylka, K. A. (2009). Pesticide multiresidue analysis in cereal grains using modified QuEChERS method combined with automated direct sample introduction GC-TOFMS and UPLC-MS/MS techniques†. *J. Agric. Food Chem.*, 58(10):5959–5972. <http://pubs.acs.org/doi/abs/10.1021/jf9029892>

- Matamoros, V., Jover, E., and Bayona, J. M. (2009). Part-per-Trillion determination of pharmaceuticals, pesticides, and related organic contaminants in river water by Solid-Phase extraction followed by comprehensive Two-Dimensional gas chromatography Time-of-Flight mass spectrometry. *Anal. Chem.*, 82(2):699–706. <http://pubs.acs.org/doi/abs/10.1021/ac902340e>
- McGorin, R. J. (2009). One hundred years of progress in food analysis. *J. Agric. Food Chem.*, 57(18):8076–8088. <http://pubs.acs.org/doi/abs/10.1021/jf900189s>
- Melbye, A. G., Brakstad, O. G., Hokstad, J. N., Gregersen, I. K., Hansen, B. H., Booth, A. M., Rowland, S. J., and Tollefsen, K. E. (2009). Chemical and toxicological characterization of an unresolved complex mixture-rich biodegraded crude oil. *Environmental Toxicology and Chemistry*, 28(9):1815–1824. <http://onlinelibrary.wiley.com/doi/10.1897/08-545.1/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Muscalu, A. M., Reiner, E. J., Liss, S. N., and Chen, T. (2009). Determination of polychlorinated biphenyls, organochlorine pesticides, chlorobenzenes in sludge and sediment samples by GCx GC $\mu$ ECD. *International Journal of Environmental Analytical Chemistry*, 90(1):1–13. <http://www.tandfonline.com/doi/abs/10.1080/03067310903229919>
- Ratel, J. and Engel, E. (2009). Determination of benzenic and halogenated volatile organic compounds in animal-derived food products by one-dimensional and comprehensive two-dimensional gas chromatography-mass spectrometry. *Journal of Chromatography A*, 1216(45):7889–7898. <http://www.sciencedirect.com/science/article/pii/S002196730901348X>
- Rega, B., Guerard, A., Delarue, J., Maire, M., and Giampaoli, P. (2009). On-line dynamic HS-SPME for monitoring endogenous aroma compounds released during the baking of a model cake. *Food Chemistry*, 112(1):9–17. <http://www.sciencedirect.com/science/article/pii/S0308814608005840>
- Rochat, S., Egger, J., and Chaintreau, A. (2009). Strategy for the identification of key odorants: Application to shrimp aroma. *Journal of Chromatography A*, 1216(36):6424–6432. <http://www.sciencedirect.com/science/article/pii/S0021967309010553>
- Schuman, M. C., Heinzl, N., Gaquerel, E., Svatos, A., and Baldwin, I. T. (2009). Polymorphism in jasmonate signaling partially accounts for the variety of volatiles produced by *Nicotiana attenuata* plants in a native population. *New Phytologist*, 183(4):1134–1148. <http://onlinelibrary.wiley.com/doi/10.1111/j.1469-8137.2009.02894.x/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Silva, A. I., Pereira, H. M. G., Casilli, A., Conceição, F. C., and Aquino Neto, F. R. (2009). Analytical challenges in doping control: Comprehensive two-dimensional gas chromatography with time of flight mass spectrometry, a promising option. *Journal of Chromatography A*, 1216(14):2913–2922. <http://www.sciencedirect.com/science/article/pii/S0021967308017627>
- Skov, T., Hoggard, J. C., Bro, R., and Synovec, R. E. (2009). Handling within run retention time shifts in two-dimensional chromatography data using shift correction and modeling. *Journal of Chromatography A*, 1216(18):4020–4029. <http://www.sciencedirect.com/science/article/pii/S0021967309002854>
- Vonderheide, A. P. (2009). A review of the challenges in the chemical analysis of the polybrominated diphenyl ethers. *Microchemical Journal*, 92(1):49–57. <http://www.sciencedirect.com/science/article/pii/S0026265X08001616>
- Windt, M., Meier, D., Marsman, J. H., Heeres, H. J., and de Koning, S. (2009). Micro-pyrolysis of technical lignins in a new modular rig and product analysis by GC-MS/FID and GCxGC-TOFMS/FID. *Journal of Analytical and Applied Pyrolysis*, 85(1-2):38–46. <http://www.sciencedirect.com/science/article/pii/S0165237008001824>
- Wu, Z., Huang, Z., Lehmann, R., Zhao, C., and Xu, G. (2009). The application of Chromatography-Mass spectrometry: Methods to metabonomics. *Chromatographia*, 69(0):23–32. <http://www.springerlink.com/content/y3n3hx64123377k5/>
- Yang, S., Sadilek, M., Synovec, R. E., and Lidstrom, M. E. (2009). Liquid chromatography-tandem quadrupole mass spectrometry and comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry measurement of targeted metabolites of *Methylobacterium extorquens* AM1 grown on two different carbon sources. *Journal of chromatography. A*, 1216(15):3280–3289. <http://www.sciencedirect.com/science/article/pii/S0021967309002544>

## 2008

---

- Adam, F., Bertoncini, F., Coupard, V., Charon, N., Thiébaud, D., Espinat, D., and Hennion, M.-C. (2008a). Using comprehensive two-dimensional gas chromatography for the analysis of oxygenates in middle distillates. *Journal of Chromatography A*, 1186(1-2):236–244. <http://www.sciencedirect.com/science/article/pii/S0021967307021292>
- Adam, F., Vendevre, C., Bertoncini, F., Thiébaud, D., Espinat, D., and Hennion, M.-C. (2008b). Comprehensive two-dimensional gas chromatography for enhanced analysis of naphthas: New column combination involving permethylated cyclodextrin in the second dimension. *Journal of Chromatography A*, 1178(1-2):171–177. <http://www.sciencedirect.com/science/article/pii/S0021967307018638>
- Akoto, L., Stellaard, F., Irth, H., Vreuls, R. J. J., and Pel, R. (2008). Improved fatty acid detection in micro-algae and aquatic meiofauna species using a direct thermal desorption interface combined with comprehensive gas chromatography“time-of-flight mass spectrometry. *Journal of Chromatography A*, 1186(1-2):254–261. <http://www.sciencedirect.com/science/article/pii/S0021967308002227>
- Amador-Muñoz, O. and Marriott, P. J. (2008). Quantification in comprehensive two-dimensional gas chromatography and a model of quantification based on selected summed modulated peaks. *Journal of Chromatography A*, 1184(1-2):323–340. <http://www.sciencedirect.com/science/article/pii/S0021967307017827>

- Aura, A.-M., Mattila, I., Seppänen-Laakso, T., Miettinen, J., Oksman-Caldentey, K.-M., and Orei, M. (2008). Microbial metabolism of catechin stereoisomers by human faecal microbiota: Comparison of targeted analysis and a non-targeted metabolomics method. *Phytochemistry Letters*, 1(1):18–22. <http://www.sciencedirect.com/science/article/pii/S1874390007000055>
- Bedair, M. and Sumner, L. W. (2008). Current and emerging mass-spectrometry technologies for metabolomics. *TrAC Trends in Analytical Chemistry*, 27(3):238–250. <http://www.sciencedirect.com/science/article/pii/S0165993608000071>
- Betancourt, S. S., Ventura, G. T., Pomerantz, A. E., Vilorio, O., Dubost, F. X., Zuo, J., Monson, G., Bustamante, D., Purcell, J. M., Nelson, R. K., Rodgers, R. P., Reddy, C. M., Marshall, A. G., and Mullins, O. C. (2008). Nanoaggregates of asphaltenes in a reservoir crude oil and reservoir connectivity. *Energy Fuels*, 23(3):1178–1188. <http://pubs.acs.org/doi/abs/10.1021/ef800598a>
- Booth, A. M., Scarlett, A. G., Lewis, C. A., Belt, S. T., and Rowland, S. J. (2008). Unresolved complex mixtures (UCMs) of aromatic hydrocarbons: Branched alkyl indanes and branched alkyl tetralins are present in UCMs and accumulated by and toxic to, the mussel *mytilus edulis*. *Environ. Sci. Technol.*, 42(21):8122–8126. <http://pubs.acs.org/doi/abs/10.1021/es801601x>
- Cajka, T., Hajslova, J., Lacina, O., Mastovska, K., and Lehotay, S. J. (2008). Rapid analysis of multiple pesticide residues in fruit-based baby food using programmed temperature vaporiser injection-low-pressure gas chromatography-high-resolution time-of-flight mass spectrometry. *Journal of Chromatography A*, 1186(1-2):281–294. <http://www.sciencedirect.com/science/article/pii/S0021967307021176>
- Cardeal, Z. L., de Souza, P. P., Silva, and Marriott, P. J. (2008). Comprehensive two-dimensional gas chromatography for fingerprint pattern recognition in cachaça production. *Talanta*, 74(4):793–799. <http://www.sciencedirect.com/science/article/pii/S0039914007004912>
- Cochran, J. (2008). Evaluation of comprehensive two-dimensional gas chromatography - time-of-flight mass spectrometry for the determination of pesticides in tobacco. *Journal of Chromatography A*, 1186(1-2):202–210. <http://www.sciencedirect.com/science/article/pii/S0021967308000812>
- Das, K. C. and Xia, K. (2008). Transformation of 4-nonylphenol isomers during biosolids composting. *Chemosphere*, 70(5):761–768. <http://www.sciencedirect.com/science/article/pii/S0045653507009174>
- de Koning, S., Kaal, E., Janssen, H.-G., van Platerink, C., and Brinkman, U. A. (2008). Characterization of olive oil volatiles by multi-step direct thermal desorption-comprehensive gas chromatography-time-of-flight mass spectrometry using a programmed temperature vaporizing injector. *Journal of Chromatography A*, 1186(1-2):228–235. <http://www.sciencedirect.com/science/article/pii/S0021967307009491>
- Dorman, F. L., Overton, E. B., Whiting, J. J., Cochran, J. W., and Gardea-Torresdey, J. (2008a). Gas chromatography. *Anal. Chem.*, 80(12):4487–4497. <http://pubs.acs.org/doi/abs/10.1021/ac800714x>
- Dorman, F. L., Schettler, P. D., Vogt, L. A., and Cochran, J. W. (2008b). Using computer modeling to predict and optimize separations for comprehensive two-dimensional gas chromatography. *Journal of Chromatography A*, 1186(1-2):196–201. <http://www.sciencedirect.com/science/article/pii/S0021967307022261>
- Dunn, W. B., Broadhurst, D., Ellis, D. I., Brown, M., Halsall, A., O'Hagan, Steven, Spasic, I., Tseng, A., and Kell, D. B. (2008). A GC-TOF-MS study of the stability of serum and urine metabolomes during the UK biobank sample collection and preparation protocols. *International Journal of Epidemiology*, 37(suppl 1):i23–i30. [http://ije.oxfordjournals.org/content/37/suppl\\_1/i23](http://ije.oxfordjournals.org/content/37/suppl_1/i23)
- Garcia, D. E., Baidoo, E. E., Benke, P. I., Pingitore, F., Tang, Y. J., Villa, S., and Keasling, J. D. (2008). Separation and mass spectrometry in microbial metabolomics. *Current Opinion in Microbiology*, 11(3):233–239. <http://www.sciencedirect.com/science/article/pii/S1369527408000532>
- Gomes da Silva, M. D. R., Cardeal, Z., and Marriott, P. J. (2008). Comprehensive Two-Dimensional gas chromatography: Application to aroma and essential oil analysis. 988:3–24. <http://pubs.acs.org/doi/abs/10.1021/bk-2008-0988.ch001>
- Graham, B. F., May, E. F., and Trengove, R. D. (2008). Emulsion inhibiting components in crude oils. *Energy Fuels*, 22(2):1093–1099. <http://pubs.acs.org/doi/abs/10.1021/ef700529m>
- Gröger, T., Schäffer, M., Pütz, M., Ahrens, B., Drew, K., Eschner, M., and Zimmermann, R. (2008a). Application of two-dimensional gas chromatography combined with pixel-based chemometric processing for the chemical profiling of illicit drug samples. *Journal of Chromatography A*, 1200(1):8–16. <http://www.sciencedirect.com/science/article/pii/S0021967308008297>
- Gröger, T., Welthagen, W., Mitschke, S., Schäffer, M., and Zimmermann, R. (2008b). Application of comprehensive two-dimensional gas chromatography mass spectrometry and different types of data analysis for the investigation of cigarette particulate matter. *J. Sep. Science*, 31(19):3366–3374. <http://onlinelibrary.wiley.com/doi/10.1002/jssc.200800340/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Haglund, P., Korytár, P., Danielsson, C., Diaz, J., Wiberg, K., Leonards, P., Brinkman, U. A. T., and Boer, J. (2008). GCxGC-ECD: a promising method for the determination of dioxins and dioxin-like PCBs in food and feed. *Analytical and Bioanalytical Chemistry*, 390(7):1815–1827. <http://www.springerlink.com/content/e537k836r6313165/>
- Hoggard, J. C. and Synovec, R. E. (2008). Automated resolution of nontarget analyte signals in GC x GC-TOFMS data using parallel factor analysis. *Anal. Chem.*, 80(17):6677–6688. <http://pubs.acs.org/doi/abs/10.1021/ac800624e>
- Hoh, E., Lehotay, S. J., Mastovska, K., and Huwe, J. K. (2008). Evaluation of automated direct sample introduction with comprehensive two-dimensional gas chromatography/time-of-flight mass spectrometry for the screening analysis of dioxins in fish oil. *Journal of Chromatography A*, 1201(1):69–77. <http://www.sciencedirect.com/science/article/pii/S0021967308009503>



- Huang, X. and Regnier, F. E. (2008). Differential metabolomics using stable isotope labeling and Two-Dimensional gas chromatography with Time-of-Flight mass spectrometry. *Anal. Chem.*, 80(1):107–114. <http://pubs.acs.org/doi/abs/10.1021/ac071263f>
- Jele, H. H. and Wasowicz, E. (2008). Determination of trichothecenes in wheat grain without sample cleanup using comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry. *Journal of Chromatography A*, 1215(1-2):203–207. <http://www.sciencedirect.com/science/article/pii/S0021967308019419>
- Kaal, E., de Koning, S., Brudin, S., and Janssen, H.-G. (2008). Fully automated system for the gas chromatographic characterization of polar biopolymers based on thermally assisted hydrolysis and methylation. *Journal of Chromatography A*, 1201(2):169–175. <http://www.sciencedirect.com/science/article/pii/S002196730800887X>
- Kallio, M. and Hyötyläinen, T. (2008). Simple calibration procedure for comprehensive two-dimensional gas chromatography. *Journal of Chromatography A*, 1200(2):264–267. <http://www.sciencedirect.com/science/article/pii/S0021967308010376>
- Kallio, M., Jussila, M., Raimi, P., and Hyötyläinen, T. (2008). Modified semi-rotating cryogenic modulator for comprehensive two-dimensional gas chromatography. *Analytical and Bioanalytical Chemistry*, 391(6):2357–2363. <http://www.springerlink.com/content/r51113716pp81g42/>
- Klimánková, E., Holadová, K., Hajlová, J., aĭka, T., Poustka, J., and Koudela, M. (2008). Aroma profiles of five basil (*ocimum basilicum* L.) cultivars grown under conventional and organic conditions. *Food Chemistry*, 107(1):464–472. <http://www.sciencedirect.com/science/article/pii/S0308814607007182>
- Li, M., Zhang, S., Jiang, C., Zhu, G., Fowler, M., Achal, S., Milovic, M., Robinson, R., and Larter, S. (2008). Two-dimensional gas chromatograms as fingerprints of sour gas-associated oils. *Organic Geochemistry*, 39(8):1144–1149. <http://www.sciencedirect.com/science/article/pii/S0146638008000466>
- Luedemann, A., Strassburg, K., Erban, A., and Kopka, J. (2008). TagFinder for the quantitative analysis of gas chromatography-mass spectrometry (GC-MS)-based metabolite profiling experiments. *Bioinformatics*, 24(5):732–737. <http://bioinformatics.oxfordjournals.org/content/24/5/732>
- Mao, D., Van De Weghe, H., Diels, L., De Brucker, N., Lookman, R., and Vanermen, G. (2008). High-performance liquid chromatography fractionation using a silver-modified column followed by two-dimensional comprehensive gas chromatography for detailed group-type characterization of oils and oil pollutions. *Journal of Chromatography A*, 1179(1):33–40. <http://www.sciencedirect.com/science/article/pii/S0021967307017311>
- Marsman, J. H., Wildschut, J., Evers, P., de Koning, S., and Heeres, H. J. (2008). Identification and classification of components in flash pyrolysis oil and hydrodeoxygenated oils by two-dimensional gas chromatography and time-of-flight mass spectrometry. *Journal of Chromatography A*, 1188(1):17–25. <http://www.sciencedirect.com/science/article/pii/S0021967308002719>
- Mateus, E. P., Gomes da Silva, M. D. R., Ribeiro, A. B., and Marriott, P. J. (2008). Qualitative mass spectrometric analysis of the volatile fraction of creosote-treated railway wood sleepers by using comprehensive two-dimensional gas chromatography. *Journal of Chromatography A*, 1178(1-2):215–222. <http://www.sciencedirect.com/science/article/pii/S0021967307020730>
- Mitrevski, B. S., Brenna, J. T., Zhang, Y., and Marriott, P. J. (2008). Application of comprehensive two-dimensional gas chromatography to sterols analysis. *Journal of Chromatography A*, 1214(1-2):134–142. <http://www.sciencedirect.com/science/article/pii/S0021967308018037>
- Mohler, R. E., Tu, B. P., Dombek, K. M., Hoggard, J. C., Young, E. T., and Synovec, R. E. (2008). Identification and evaluation of cycling yeast metabolites in two-dimensional comprehensive gas chromatography-time-of-flight-mass spectrometry data. *Journal of Chromatography A*, 1186(1-2):401–411. <http://www.sciencedirect.com/science/article/pii/S0021967307018092>
- Mondello, L., Tranchida, P. Q., Dugo, P., and Dugo, G. (2008). Comprehensive two-dimensional gas chromatography-mass spectrometry: A review. *Mass Spectrom. Rev.*, 27(2):101–124. <http://onlinelibrary.wiley.com/doi/10.1002/mas.20158/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- O'Neal, K. L., Fang, H., Guy, Y., Hamsher, A., Hong, L., Li, L., Wang, M., Xu, H., Xu, X., Yang, Y., and Weber, S. G. (2008). Pittcon provides a platform for interesting, important and novel work. *TrAC Trends in Analytical Chemistry*, 27(4):274–283. <http://www.sciencedirect.com/science/article/pii/S0165993608000836>
- Orei, M., Simell, S., Sysi-Aho, M., Nántö-Salonen, K., Seppänen-Laakso, T., Parikka, V., Katajamaa, M., Hekkala, A., Mattila, I., Keskinen, P., Yetukuri, L., Reinikainen, A., Lähde, J., Suortti, T., Hakalax, J., Simell, T., Hyöty, H., Veijola, R., Ilonen, J., Laheesmaa, R., Knip, M., and Simell, O. (2008). Dysregulation of lipid and amino acid metabolism precedes islet autoimmunity in children who later progress to type 1 diabetes. *The Journal of Experimental Medicine*, 205(13):2975–2984. <http://jem.rupress.org/content/205/13/2975>
- Pangallo, K., Nelson, R. K., Teuten, E. L., Pedler, B. E., and Reddy, C. M. (2008). Expanding the range of halogenated 1'-methyl-1,2'-bipyrroles (MBPs) using GC/ECNI-MS and GCxGC/TOF-MS. *Chemosphere*, 71(8):1557–1565. <http://www.sciencedirect.com/science/article/pii/S0045653507014853>
- Pasikanti, K. K., Ho, P. C., and Chan, E. C. Y. (2008). Gas chromatography/mass spectrometry in metabolic profiling of biological fluids. *Journal of Chromatography B*, 871(2):202–211. <http://www.sciencedirect.com/science/article/pii/S1570023208002857>
- Peters, R., Tonoli, D., van Duin, M., Mommers, J., Mengerink, Y., Wilbers, A. T. M., van Benthem, R., de Koster, C., Schoenmakers, P. J., and van der Wal, S. (2008). Low-molecular-weight model study of peroxide cross-linking of ethylene-propylene (-diene) rubber using gas chromatography and mass spectrometry. *Journal of Chromatography A*, 1201(2):141–150. <http://www.sciencedirect.com/science/article/pii/S0021967308005888>

- Qiu, Y., Lu, X., Pang, T., Ma, C., Li, X., and Xu, G. (2008). Determination of radix ginseng volatile oils at different ages by comprehensive two-dimensional gas chromatography/time-of-flight mass spectrometry. *J. Sep. Science*, 31(19):3451–3457. <http://onlinelibrary.wiley.com/doi/10.1002/jssc.200800253/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Ralston-Hooper, K., Hopf, A., Oh, C., Zhang, X., Adamec, J., and Sepúlveda, M. S. (2008). Development of GCxGC/TOF-MS metabolomics for use in ecotoxicological studies with invertebrates. *Aquatic Toxicology*, 88(1):48–52. <http://www.sciencedirect.com/science/article/pii/S0166445X08000854>
- Ryona, I., Pan, B. S., Intrigliolo, D. S., Lakso, A. N., and Sacks, G. L. (2008). Effects of cluster light exposure on 3-Isobutyl-2-methoxypyrazine accumulation and degradation patterns in red wine grapes (*vitis vinifera* L. cv. cabernet franc). *J. Agric. Food Chem.*, 56(22):10838–10846. <http://pubs.acs.org/doi/abs/10.1021/jf801877y>
- Scarlett, A., Rowland, S. J., Galloway, T. S., Lewis, A. C., and Booth, A. M. (2008). Chronic sublethal effects associated with branched alkylbenzenes bioaccumulated by mussels. *Environmental Toxicology and Chemistry*, 27(3):561–567. <http://onlinelibrary.wiley.com/doi/10.1897/07-347.1/abstract?systemMessage=Wiley+Online+Library+will+be+disrupted+on+27+October+from+10%3A00-12%3A00+BST+%2805%3A00-07%3A00+EDT%29+for+essential+maintenance>
- Schurek, J., Portolés, T., Hajslova, J., Riddellova, K., and Hernández, F. (2008). Application of head-space solid-phase microextraction coupled to comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry for the determination of multiple pesticide residues in tea samples. *Analytica Chimica Acta*, 611(2):163–172. <http://www.sciencedirect.com/science/article/pii/S0003267008000755>
- Stepan, R., Cuhra, P., and Barsova, S. (2008). Comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometric detection for the determination of anabolic steroids and related compounds in nutritional supplements. *Food Additives & Contaminants: Part A*, 25(5):557–565. <http://www.tandfonline.com/doi/abs/10.1080/02652030701609228>
- Tran, T. C. and Marriott, P. J. (2008). Comprehensive two-dimensional gas chromatography - time-of-flight mass spectrometry and simultaneous electron capture detection/nitrogen phosphorous detection for incense analysis. *Atmospheric Environment*, 42(32):7360–7372. <http://www.sciencedirect.com/science/article/pii/S1352231008006237>
- van der Lee, M. K., van der Weg, G., Traag, W. A., and Mol, H. G. J. (2008). Qualitative screening and quantitative determination of pesticides and contaminants in animal feed using comprehensive two-dimensional gas chromatography with time-of-flight mass spectrometry. *Journal of Chromatography A*, 1186(1-2):325–339. <http://www.sciencedirect.com/science/article/pii/S0021967307019140>
- van Leeuwen, S. P. J. and de Boer, J. (2008). Advances in the gas chromatographic determination of persistent organic pollutants in the aquatic environment. *Journal of Chromatography A*, 1186(1-2):161–182. <http://www.sciencedirect.com/science/article/pii/S0021967308001337>
- Ventura, G. T., Kenig, F., Reddy, C. M., Frysinger, G. S., Nelson, R. K., Mooy, B. V., and Gaines, R. B. (2008). Analysis of unresolved complex mixtures of hydrocarbons extracted from late archean sediments by comprehensive two-dimensional gas chromatography (GC<sub>A</sub>—GC). *Organic Geochemistry*, 39(7):846–867. <http://www.sciencedirect.com/science/article/pii/S014663800800079X>





**LECO Corporation**

3000 Lakeview Avenue • St. Joseph, MI 49085 • Phone: 800-292-6141 • Fax: 269-982-8977  
info@leco.com • www.leco.com • ISO-9001:2000 • No. FM 24045 • LECO is a registered trademark of LECO Corporation.